

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
**WO 01/91726 A1**

(51) International Patent Classification<sup>7</sup>: A61K 9/06, 7/48

S. [US/US]; 1521 Ide Drive, Lansdale, PA 19446 (US).  
**DORRELL, Jay** [US/US]; 1459 Woodford Drive, Wayne,  
PA 19087 (US).

(21) International Application Number: PCT/US01/18033

(22) International Filing Date: 1 June 2001 (01.06.2001)

(74) Agents: **BARRON, Alexis et al.**; Synnestvedt & Lechner  
LLP, 2600 Aramark Tower, 1101 Market Street, Philadel-  
phia, PA 19107-2950 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/209,330 2 June 2000 (02.06.2000) US  
0023891.5 29 September 2000 (29.09.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,  
SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,  
ZW.

(71) Applicant (*for all designated States except US*): **DER-  
MIK INTERNATIONAL HOLDING INC.** [US/US];  
3711 Kennett Pike, Suite 200, Greenville, DE 19807 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **MANETTA,  
Vincent, E.** [US/US]; Galderma, 5 Cedarbrook Drive,  
Cranbury, NJ 08512 (US). **MAHONEY, Linda, M.**  
[US/US]; 103 Beaufort Court, Phoenixville, PA 19460  
(US). **NORDSIEK, Michael, T.** [US/US]; 561 Bair  
Road, Berwyn, PA 19312 (US). **BALAJI, Kodumudi,**

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

[Continued on next page]

(54) Title: ACNE-TREATING COMPOSITION



(57) Abstract: A package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, the composition tending to degrade prematurely, one of the components comprising an oxidizing agent and another of the components comprising an antibiotic which is effective against acne-associated bacterial species, the components separated one from the other in the package, one component having a viscosity within about 50 % of the viscosity of the other component. In addition, a package comprising: (A) components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of the components comprising a benzoyl peroxide gel and another of the components comprising a gel of erythromycin and hydroxypropylcellulose; and (B) containers for holding the components in the package separated one from the other; the components having viscosities such that, upon the application of a uniform forces to the components, substantially equal volumes of the components are capable of being dispensed simultaneously from the containers.

WO 01/91726 A1

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

-1-

**ACNE-TREATING COMPOSITION****Field of the Invention**

The present invention relates to a pharmaceutical composition that tends to lose its effectiveness prematurely  
5 and to improved means for the effective use of such a composition. More particularly, the present invention relates to the packaging of constituents comprising such a composition and to an improved form of such composition, including, for example, a composition for topical  
10 application, such as a topical composition for the treatment of acne.

Acne is a common inflammatory disease which is very common at puberty and may continue for many years. It occurs in facial skin areas where sebaceous glands are the largest,  
15 most numerous, and most active. In its milder forms, acne is a superficial disorder which is evidenced by slight, spotty irritations and which can be treated satisfactorily by ordinary skin hygiene. However, in the more inflammatory types of acne, bacterial invasion of or about the

pilosebaceous follicles occurs and results in the formation of pustules, infected cysts, and, in extreme cases, canalizing inflamed and infected sacs appear. These lesions may become extensive and leave permanent, disfiguring scars.

5       Therapeutic methods for treating acne are designed to prevent formation of new lesions and facilitate the healing of old lesions. Treatments include the systemic and topical administration of anti-acne agents such as antibiotics or synthetic Vitamin A analogs. In most cases, systemic  
10 treatment of acne is not desirable because of the risks of adverse side effects that are experienced by the user. For this reason, topical acne treatment compositions have been preferred.

      The present invention relates to topical anti-acne  
15 compositions and packages which contain the components of the composition.

#### Reported Developments

      Topical anti-acne compositions include, for example, one or more of the following: sulfur, resorcinol, salicylic acid,  
20 benzoyl peroxide, steroids and antibiotics. Exemplary antibiotics are disclosed in U.S. Patent No. 3,969,516 (lincomycin family); BR Publication No. 1,594,314 (erythromycin); and U.S. Patent No. 3,952,099 (tetracycline). Preparations containing a peroxide are reported in U.S.  
25 Patent Nos. 3,535,422; 4,056,611; 4,387,107, and British Publication No. 1,594,314 and U.S. Patent No. 4,497,794. Antibiotic-containing compositions which also include anti-

inflammatory steroids are disclosed in U.S. Patent No. 4,132,781.

Attempts to improve the effectiveness of topical antibiotic compositions for use in the treatment of acne have taken a number of approaches. One approach is reported in U.S. Patent Nos. 3,989,815; 3,989,816, 3,991,203; 4,122,170; 4,316,893; 4,444,762; and EP 27,286, which disclose skin-penetrating vehicle compositions that reportedly increase the transdermal absorption of any physiologically active substance, including antibiotics. However, some penetrating agents are not compatible with antibiotics used to treat acne.

A further approach relates to the use of a composition which utilizes two different active components, such as erythromycin, clindamycin, Vitamin A acid or benzoyl peroxide. Compositions including mixtures of a peroxide and erythromycin are reported in British Publication No. 1,594,314. Acne-treating compositions comprising benzoyl peroxide, erythromycin, and hydroxyethylcellulose have also been disclosed in "The Formulation and Stability of Erythromycin-Benzoyl Peroxide in a Topical Gel, Vermeulen et al., *International Journal of Pharmaceutics*, 178, 137-141 (1999)". U.S. Patent No. 5,894,019 to Hess et al. discloses a topically applied pharmaceutical composition that is prepared by mixing a liquid solution of benzoyl peroxide (or other pharmaceutically active ingredient) that is dissolved in a liquid lipid with a hydrous gel that may also contain a pharmaceutically active ingredient, such as erythromycin. The gelling agent used to form the hydrous gel may include

hydroxypropylcellulose. U.S. Patents Nos. 5,466,446 to Stiefel et al, 5,733,886 to Baroody et al., and 5,767,098 to Klein et al. disclose acne-treating compositions which contain clindamycin and benzoyl peroxide.

5       The above compositions comprising a peroxide, such as benzoyl peroxide, and an antibiotic, such as erythromycin or clindamycin, may have stability limitations. When the peroxide and antibiotic are combined, the peroxide can partially or completely oxidize the antibiotic, rendering it  
10   ineffective. If the peroxide and antibiotic are provided separately, it is usually necessary to have a pharmacist combine the component containing the peroxide with the component containing the antibiotic to ensure that the composition is prepared properly, has the correct final  
15   concentrations of the peroxide and antibiotic, and to maximize the shelf life of the composition by postponing the mixing of the components until the time of sale. Another problem with these compositions is that once mixed, the oxidation reactions of the peroxide not only degrade the  
20   antibiotic, but also result in the accumulation of undesirable degradative products in the composition. An additional problem is that unused portions of certain acne-treating composition must be refrigerated to minimize degradation until use.

25       The present invention relates to improved means for using effectively anti-acne compositions that tend to lose their effectiveness prematurely and to improved forms of anti-acne compositions.

Summary of th Invention

In accordance with the present invention, there is provided a package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition  
5 that is effective in treating acne, the composition tending to degrade prematurely, one of the components comprising an oxidizing agent and another of the components comprising an antibiotic which is effective against acne-associated  
bacterial species, the components separated one from the  
10 other in the package, one component having a viscosity within about 50% of the viscosity of the other component.

In preferred embodiments, the oxidizing agent is benzoyl peroxide, the antibiotic is selected from the group consisting of erythromycin and clindamycin, and the package  
15 is a dual packet laminated foil package.

Another aspect of the present invention provides a package comprising: (A) components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of the components comprising  
20 a benzoyl peroxide gel and another of the components comprising a gel of erythromycin and hydroxypropylcellulose; and (B) containers for holding the components in the package separated one from the other; the components having viscosities such that, upon the application of a uniform  
25 force to the components, substantially equal volumes of the components are capable of being dispensed simultaneously from the containers.

Yet another aspect of the present invention is a composition for the treatment of acne comprising: (A) benzoyl

peroxide; (B) erythromycin; and (C) hydroxypropylcellulose. Examples of preferred compositions may further comprise an additional gelling agent, an agent which inhibits degradative interactions between benzoyl peroxide and erythromycin, a  
5 base which increases the pH of the composition to a value at which said composition has an acceptable viscosity, an alcohol solvent for erythromycin, and a diluent.

Yet another aspect of the present invention provides a composition for the treatment of acne comprising: (a) benzoyl  
10 peroxide; (b) erythromycin; (c) hydroxypropylcellulose; (d) a gelling agent; (e) a surface active agent; (f) a base; (g) an alcohol; and (h) water. An example of a preferred gelling agent includes a hydroxylated vinylic polymer.

An example of a preferred base is sodium hydroxide. An  
15 example of a preferred alcohol used in the composition is specially denatured ethanol. An example of a surface active agent is dioctyl sodium sulfosuccinate.

The invention includes also within its scope the provision of a package containing a first acne-treating  
20 composition and a second acne-treating composition, the compositions tending to degrade chemically upon being mixed with one another, the package comprising a first chamber containing the first composition and a second chamber containing the second composition, each of the chambers  
25 having a sealed dispensing orifice, the orifices positioned adjacent to one another.

Another aspect of the present invention provides a package containing a first acne-treating composition and a



second acne-treating composition, the compositions tending to degrade chemically upon being mixed with one another, the package comprising a first and second packet, the packets being attached to each other and each of the packets having a sealed dispensing orifice, the sealed dispensing orifices being positioned adjacent to each other so that, as the compositions are dispensed through the orifices in unsealed form, they are combined with each other.

An additional aspect of the present invention provides a package containing a first acne-treating composition and a second acne-treating composition, the compositions tending to degrade chemically upon being mixed with one another, the package comprising a first and second packet, each of the packets having a top which includes a dispensing orifice that has a tear-off tab seal, a bottom, two sides in a parallel relationship to one another, and an exterior face and an interior face, the first packet containing the first composition and the second packet containing the second composition, the interior faces of said packets adjacent to and facing one another such that the tear-off tabs can be simultaneously torn off such that the compositions are dispensed and combined with each other as they exit the packets through the dispensing orifices.

#### Description of the Drawings

Figure 1 is a plan view of a dual-packet package.

Figure 2 is a plan view of the package of Figure 1 folded about axis A-A.

Figure 3 is a cross-sectional view of a trilaminate package.

Figure 4 is an exploded view of a pump for dispensing the compositions of the present invention.

5        Figure 5 is a partial sectional view of the assembled pump depicted in Figure 4.

### Detailed Description of the Invention

The acne-treating composition of the present invention comprises a plurality of components which when combined tend  
10 to degrade prematurely. Degradation is considered premature when the components are mixed together and at least one of the components degrades at a rate faster than the rate at which it degrades (if at all) in the absence of the other component. As an example, in acne-treating compositions  
15 which contain an oxidizing agent and an antibiotic, the oxidizing agent will oxidize and thereby degrade the antibiotic at a much faster rate than the antibiotic would degrade in the absence of the oxidizing agent.

Components that degrade prematurely when combined may be  
20 used in the practice of the present invention and may be identified using techniques known in the art. Examples of such components include, but are not limited to, oxidizing agents, gelling agents, and antibiotics.

The acne-treating composition of the present invention  
25 is provided in a package which maintains the components separated from each other until the components are removed in whole or in part from the package. In preferred form, the package is designed to enable the user to extract readily

from the package amounts of components which form an effective acne-treating composition.

Use of such a package allows long-term storage of the components at room temperature without degradation. The package can be carried conveniently by the patient because the contents do not require refrigeration. This increases patient compliance with administration of the composition. In addition, because the composition is mixed at the time of application, degradation is minimized. Furthermore, the proportions of active ingredients in the composition can be optimized by exact filling of packets and by the ease with which the patient can empty the packets simultaneously such that the final composition has the desired ratios of components.

The package of the present invention may be made from any suitable material, for example, paper, plastic, or metal foil, in laminated or composite form. The material functions to protect the components from exposure to air, moisture, light and other environmental factors which could change the physical and/or chemical nature of the components and to keep the components from contact with each other.

In preferred form, the package of the present invention includes chambers which maintain the components of the composition in separated form and which accommodate amounts of the components such that, when the components are removed from their chambers and brought together, they form a single effective acne-treating dose of the composition. For example, in preferred embodiments, the package contains a sufficient amount of the composition to apply a thin layer of

the composition to the entire surface of the face (with the exception of the eyes and mouth) and sufficient amounts of the pharmaceutically active ingredients to treat effectively any acne present in the areas to which the composition is  
5 applied.

It is believed that the present invention will be used widely in the packaging of an acne-treating composition comprising a component which includes an oxidizing agent and a component which includes an antibiotic which is effective  
10 in treating an acne condition. As mentioned above, the oxidizing agent in such a composition tends to oxidize and thereby degrade the antibiotic when the two components are brought together. The present invention may also be used in acne-treating compositions which include retinoic acid.

15 A particularly preferred embodiment of the package of the present invention is shown in Figure 1, in which there is illustrated a dual-packet, laminated metal foil package 1 comprising a first packet 5 which includes a first chamber 2 containing a first component of the pharmaceutical  
20 composition and a second packet 7 which includes a second chamber 3 containing a second component of the composition. Each of the chambers 2 and 3 has a sealed dispensing orifice 9 and 9' with tear-off tabs 11 and 11'. Each packet has a top portion 13 and 13', a bottom portion 15 and 15', a  
25 commonly shared side 17 and a second side 19 and 19' in parallel relationship to each other, and an interior face 21 and 21' and an exterior face (not shown in Figure 1) in opposed relationship to each other. The first chamber 2 and second chamber 3 may be of equal or different size.

Figure 2 shows the package of Figure 1 in a form in which the packets have been folded about axis A and the interior faces 21 and 21' of the packet are in apposition to one another and the exterior faces 23 and 23' (not shown) face outwardly. The tear-off tabs 11 and 11' of the packets are superimposed upon one another. This permits them to be removed simultaneously. After removal of the tabs, the components in chambers 2 and 3 can be dispensed and combined with each other as they exit packets 5 and 7 through the dispensing orifices 9 and 9'.

In preferred embodiments, the packages are comprised of two packets made of a multi-layered laminated foil. Preferably the laminate is a five-layer design consisting of three main barrier layers and two adhesive layers which lie between the barrier layers and hold the layers together. In preferred embodiments, the packets have a first innermost layer comprised of low density polyethylene. This layer is in contact with the composition. The second layer is an adhesive layer comprised of ethylene-acrylic acid copolymer which binds the first layer to the third layer. The third layer is a barrier layer made of aluminum foil. The fourth layer is comprised of ethylene-acrylic acid copolymer which serves as an adhesive layer binding the third layer to the fifth layer. The fifth and final layer is comprised of polyester which provides durability and tear- and puncture-resistance to the package. A preferred laminate for use in the present invention is available from Reynolds Metals Co.

For exemplary purposes, the use of the package of Figures 1 and 2 is described in connection with an acne-

treating composition comprising an oxidizing agent and an antibiotic. Accordingly, one of the chambers of the package contains the oxidizing agent and the other chamber contains the antibiotic. To extract the oxidizing agent and

5 antibiotic from the package, both packets thereof are opened simultaneously by tearing off their respective tear-off tabs. The packets are then squeezed by the user until the contents of the chambers are emptied substantially. The contents typically exit their respective orifices in the form of a

10 ribbon. The oxidizing agent and the antibiotic component can be mixed conveniently to form the acne-treating composition by the user's dispensing the contents into the palm of one hand and using the fingers of the other hand to mix the components together by use of a swirling motion. The

15 composition is then applied to the site or sites in need of treatment.

The package of Figures 1 and 2 can be characterized as a single-dose package which contains a sufficient amount of each component to form a volume of composition which can be

20 applied as a thin layer to the entire surface of the face (with the exception of the eyes and mouth) with a minimal amount of the composition left over. Any remaining composition may be discarded. Because of the relatively small amounts of components in the package, the package can

25 be considered also to be a "single-use" package which is thrown away after its contents are emptied.

Figure 3 illustrates a cross-sectional view of another package embodiment in which there is illustrated a "trilaminate" package 26 comprising a first packet 27 and a

second packet 29, having a common internal barrier layer 31, preferably a single layer of aluminum foil or other suitable material known in the art. This internal barrier layer 31 is in contact with the first 32 and second 33 compositions within the first 27 and second 29 packets. The internal barrier layer 31 is sealed along its perimeter to a first exterior barrier layer 34, and a second exterior barrier layer 36. Both the first 34 and second 36 exterior barrier layers are preferably comprised of a single layer, of aluminum foil or other suitable material known in the art.

A "multiple use" package is capable of holding relatively large amounts of components for use in forming the treating composition. The multiple-use package is similar to the single-use package in that the components of the acne-treating composition are kept separate from one another and mixed when the product is dispensed. In preferred embodiments, the multiple-use package may include a dose-metering pump which dispenses an equal amount of both components of the acne-treatment composition.

Figure 4 shows an exploded side view of a pump 50 for dispensing a multi-component fluid formulation, for example, a topical anti-acne composition. The preferred pump embodiment shown is designed to accommodate a two-component formulation, although it is understood that the design is not limited to use with formulations having only two components.

Pump 50 is suitable for use with lotions, creams, oils, gels and liquids and is designed to keep the components of the formulation separate while they are stored in the pump. Mixing of the components occurs only after they have been

dispensed from the pump immediately before use. This feature is useful to extend the effective life of formulations, such as the anti-acne composition, which tend to degrade once the components are combined. The pump is also designed to

5 dispense the components of the formulation in the proper relative amounts which provide maximum effectiveness of the formulation.

As seen in Figure 4, pump 50 comprises a bottle 52 having two reservoirs 54a and 54b separated from one another

10 by a bulkhead 56. The reservoirs hold the different components of the formulation and are attached to a common neck 58 which provides for attachment of the other pump components, described below, to the bottle 52. Neck 58 surrounds and defines two openings 60a and 60b in reservoirs

15 54a and 54b respectively, the openings permitting the components of the formulation to be loaded into the reservoirs. A liner 62 engages neck 58 and seals openings 60a and 60b to prevent premature mixing of the components within the pump.

20 Liner 62 has two through openings 64a and 64b through which dip tubes 66a and 66b extend into reservoirs 54a and 54b respectively. Dip tubes 66a and 66b provide conduits for drawing the formulation components from the reservoirs 54a and 54b to be dispensed by the pump. Each dip tube is

25 connected to a respective valve seat 68a or 68b, the valve seats being attached to a receptacle 70 which sits atop neck 58.

As best seen in Figure 5, receptacle 70 has two cylinders 72a and 72b which are in fluid communication with



dip tubes 66a and 66b through valve seats 68a and 68b respectively. Housed in each cylinder is a respective ball 74a or 74b. Balls 74a and 74b are preferably stainless steel and are sized to seat in respective valve seats 68a and 68b to form a pair of check valves 76a and 76b which allow components of the formulation to flow in one direction only, from the respective reservoirs 54a and 54b through the dip tubes 66a and 66b and into cylinders 72a and 72b as described in detail below.

Cylinders 72a and 72b each receive a respective piston 78a and 78b. The pistons are sized to sealingly interfit slidably within their respective cylinders and provide suction and pumping action when they are reciprocated within the cylinders during pump actuation. Pistons 78a and 78b have respective piston bores 80a and 80b which provide fluid communication between their respective cylinders 72a and 72b and a spout 82 attached to the tops of the pistons. Pistons 78a and 78b are each biased to a position away from check valves 76a and 76b by respective springs 84a and 84b housed within cylinders 72a and 72b between the valve seats 68a and 68b and the pistons 78a and 78b as shown.

As seen in Figure 5, spout 82 has ducts 86a and 86b which are in fluid communication with piston bores 80a and 80b of pistons 78a and 78b respectively. A second set of valve seats 88a and 88b are provided at the tops of piston bores 80a and 80b. A second set of balls 90a and 90b are positioned within ducts 86a and 86b respectively, the balls being sized to seat on respective valve seats 88a and 88b and form a second set of check valves 92a and 92b. Preferably,

balls 90a and 90b are made of stainless steel. Ducts 86a and 86b are in fluid communication with respective dispensing nozzles 94a and 94b from which the components held in reservoirs 54a and 54b are dispensed to be mixed outside of  
5 pump 50.

As seen in Figures 4 and 5, most of the pump components including the spout 82, pistons 78a and 78b, check valves 76a, 76b, 92a and 92b, springs 84a and 84b, and receptacle 70 are housed within a retainer 96. Retainer 96 has a lower  
10 portion 98 adapted to engage neck 58 and retain the pump components to the bottle 52. Attachment of the retainer is preferably by an interference fit between the retainer inner surface 100 and raised ribs 102 surrounding neck 58. An over cap 104 fits over the retainer 96 and protects the moving  
15 parts of the pump against unintentional pump actuation.

Pump 50 is preferably made from various types of resilient, flexible plastic materials, such as polyethylene and polypropylene to cite examples. Plastics are advantageous because they are inexpensive, easily molded,  
20 substantially unbreakable, and chemically inert and, thus, will not affect the components of the formulation. Parts of the pump, such as the springs for biasing the pistons and the balls for the check valves, for which plastic is not suited are preferably made of stainless steel. Stainless steel  
25 provides a robust, durable material which also is substantially chemically inert and, thus, suitable for pump parts which will come into contact with the components of the formulation during pump operation.

Operation of the dual dispensing pump 50 is described with reference to Figure 5. In operation, the user removes the over cap 104 exposing spout 82. With the pump 50 in an upright position, the user manually depresses spout 82. This  
5 causes the spout and attached pistons 78a and 78b to move downwardly with respect to the retainer 96, check valves 76a and 76b, liner 62, and bottle 52.

Pistons 78a and 78b move within their respective cylinders 72a and 72b compressing the formulation components  
10 within the cylinders as well as springs 84a and 84b. Balls 74a and 74b are firmly seated in valve seats 68a and 68b, operating as check valves to prevent the formulation components in the cylinders from being forced into respective reservoirs 54a and 54b. The formulation components are  
15 forced upwardly through the piston bores 80a and 80b in each piston respectively, the hydraulic pressure within each cylinder unseating balls 90a and 90b from valve seats 88a and 88b at the tops of the piston bores 80a and 80b. The formulation components are forced from the cylinders through  
20 the piston bores into ducts 86a and 86b in the spout, the components being dispensed from dispensing nozzles 94a and 94b in the ducts and into the hand of the user who mixes the components to form the complete formulation, ready for use.

Upon release of spout 82, springs 84a and 84b push the  
25 pistons 78a, 78b to their biased position spaced away from check valves 76a and 76b. Upward motion of the pistons creates a suction within the cylinders 72a and 72b and the piston bores 80a and 80b. The suction seats balls 90a and 90b on valve seats 88a and 88b to prevent formulation

components in ducts 86a and 86b from being drawn back into the piston bores 80a and 80b. Simultaneously, the suction causes balls 74a and 74b to lift off their respective seats 68a and 68b as the formulation components are drawn up  
5 through dip tubes 66a and 66b from reservoirs 54a and 54b and into cylinders 72a and 72b. The pump is then ready to dispense the formulation components on the next actuation.

As explained above, each component of the formulation is pumped from its reservoir by its own piston-cylinder  
10 combination. The amount of a particular component dispensed from the pump is proportional to the volume of the cylinder swept by its piston, which pumps that particular component. Since the strokes of both pistons are equal, the relative amount of component dispensed from one reservoir in relation  
15 to the other reservoir is determined by the ratio of the cylinder diameters. Cylinders having equal diameters will have equal swept volumes and, thus, dispense equal amounts of formulation component from each reservoir. If the diameter of one of the cylinders is made smaller relative to the other  
20 cylinder, then the piston associated with the smaller cylinder will sweep a smaller volume, and a smaller amount of component will be dispensed from the respective reservoir. Thus, the pump can be designed to automatically dispense formulation components in the proper relative proportions for  
25 maximum effectiveness simply by having cylinders with different diameters, the ratio of the diameters being proportional to the desired ratio of formulation components to be dispensed.

The package embodiments of the present invention are designed to be used by the lay person who has an acne condition or a lay person who applies the composition to the individual with the condition. Accordingly, the packages are  
5 designed so that the user is able to dispense on a consistent basis effective amounts of the active ingredients comprising the pharmaceutical composition. Thus, the need for a pharmacist to prepare the composition is eliminated. The specific amount of each of the active ingredients present in  
10 the package will depend on the identity of the active ingredient, taking into account its known effect on treatment of the acne condition and the proportion of the ingredients that are desired in the composition that is formed by mixing the active ingredients. Also, the package may contain an  
15 optimal dosage of the active ingredients of the composition by including in each packet the appropriate volume and concentration of the active ingredient. For example, in the use of a peroxide/antibiotic composition, it may be desired that the peroxide comprises a greater proportion of the  
20 composition than the antibiotic.

It will be appreciated that the packets of the single-dose package, as described herein, contain relatively small volumes of the components from which the composition is formulated. For example, an embodiment of such a packet  
25 contains about 0.425 g of one component and about 0.425 g of a second component. The dispensed components are within plus or minus 0.025 g of the 0.425 g in the packet, allowing for about a 6% tolerance. The filling of the packets with effective dosage amounts of the components is essential to

providing an effective acne-treating composition. This is difficult to accomplish with components that are in a form of the type described herein, for example, in gel form. In order to fill the packets with the accurate dosages of components, it has been found that it is important to rid the components of air trapped within the components. To avoid trapped air, the components are formulated preferably using a process which avoids the introduction of air into the components and removes trapped air from the components. This is accomplished by preparing the components under a vacuum, for example, about -15 to about -30 psi. By substantially eliminating trapped air from the components, it is possible to fill the packets or other appropriate containers with a high degree of accuracy. This, in turn, allows the lay person to prepare the acne-treating composition within the desired concentration parameters.

The form of the composition that is applied to the skin should be such that it adheres to the skin for a sufficient period of time to allow the active ingredients to act on the affected area. As an example, for embodiments which include a peroxide and an antibiotic, the composition will be in a form which adheres to the skin for a sufficient period of time for the peroxide to inactivate extracellular lipases and inhibit the formation of free fatty acids in the skin and for the antibiotic to reduce the concentration of acne-associated bacteria in the affected area. For this purpose, the acne-treating composition should have a viscosity such that the composition adheres to the skin. This requires that, when the components that make up the composition are combined

together, they form a composition which has a viscosity such that the composition adheres to the skin. The forms of the components and the viscosities thereof can be different and vary substantially, provided that they can be combined to  
5 form a composition having the desired viscosity characteristics. For example, the components may be in forms ranging from liquids to solids. In preferred embodiments, as described in detail below, the component is in a form which has properties intermediate the liquid and solid states.  
10 Examples of such forms include gels, lotions, creams, salves, and ointments. In a particularly preferred embodiment, the components are in gel form.

The package of the present invention, as exemplified in Figures 1 and 2, includes the separated components in dosages  
15 which are effective in forming, upon mixing, an effective acne-treating composition. In preferred form, the package is designed and the components are formulated to enable the user to dispense on a consistent basis substantially all of the contents of the package. Accordingly, the contents are made  
20 available for combining into an acne-treating composition in which the components are present in the desired dosage. The preferred means of accomplishing this is to formulate components which have viscosities that enable the user to dispense the components from the package without undue  
25 effort. It is preferred also that the components be in the same form. For example, if one component is in gel form, it is preferred that the other component be in gel form. In addition, it is preferred that the components have similar viscosities. The term "similar viscosity" means that when a

uniform "dispensing" force is applied to the separated components, substantially equal amounts of the components are dispensed simultaneously from their respective chambers. In a particularly preferred form, the components have

5 viscosities, as measured by 1V-101 Brookfield Helipath T-F @ 1.5 rpm that are within about 50%, preferably within about 25% of each other.

Standard procedures known in the art may be used to determine viscosity. To perform the viscosity test, a test  
10 sample was transferred to a 1lb jar (89 mm in diameter and 92 mm in height) while tapping on the jar to eliminate air. After the transfer, the jar was left to sit for one hour at 25.0°C. A Brookfield Model LV series rotational viscometer equipped with a helipath and a T-F spindle was used to  
15 measure the viscosity of the sample. The spindle of the viscometer was placed in the sample and the height of the spindle was adjusted so that the crosspiece of the spindle was covered by about 1/4" of the sample. The rotational speed of the spindle was set to 1.5 rpm. The helipath and  
20 the viscometer were simultaneously started. After one minute, the needle lock was engaged. When the needle was displayed in the window, the viscometer was shut off. A reading was taken and recorded on the 0-100 scale. Additional tests may be performed to obtain additional  
25 readings. If additional tests are performed, the viscometer rotation should be resumed and at least one rotation allowed prior to the release of the needle lock. Timing of the additional sample readings is to be commenced from the time of the release of the needle lock.



In particularly preferred embodiments, each of the components is a gel which has a viscosity of about 200,000 cps to about 500,000 cps.

There follows a description of exemplary ingredients comprising the acne-treating composition of the present invention, including a description of exemplary pharmaceutically active ingredients and exemplary ingredients that are combined therewith to place the composition in a suitable form for application to the face and for dispensing the components of the composition from the package of the present invention.

In the discussion which follows, the amount of an ingredient in the acne-treating composition is expressed in weight percent based on the total weight of the composition (% w/w - composition). The amount of an ingredient in one of the components is expressed in weight percent based on the total weight of the component (% w/w - component).

The acne-treating composition includes an antibiotic which is effective against acne-associated bacterial species. An antibiotic is effective against acne-associated bacterial species if it kills or inhibits the growth of a bacterial population. The term "acne-associated bacterial species" refers to bacterial species found at the site of an acne lesion, in particular *Propionibacterium acnes* (*P. acnes*). Exemplary antibiotics include lincomycin, clindamycin (a lincomycin derivative), and erythromycin and their pharmaceutically acceptable salts and esters such as their hydrochlorides and phosphates.

Lincomycin is a derivative of the amino acid trans-L-4- $\alpha$ -propyl-hydrinic acid coupled to a derivative of an octose substituted by a methylmercaptyl group. Lincomycin antibiotics are described in U.S. Patent Nos. 3,475,407;  
5 3,509,127; 3,544,551 and 3,513,155.

Clindamycin is the 7-deoxy,7-chloro derivative of lincomycin, and is otherwise known as methyl 7-chloro-6,7,8,trideoxy-6-[[ (1-methyl-4-propyl-2-pyrrolidinyl) carbonyl]amino]-1-thio-L-threo- $\alpha$ -D-galacto-octopyranoside.  
10 In the use of clindamycin as the antibiotic, it is recommended that there be used clindamycin phosphate which is converted *in vivo* to active clindamycin.

A preferred antibiotic for use in the present invention is erythromycin, which is a macrolide antibiotic produced  
15 from a strain of *Saccharopolyspora erythraea* (formerly *Streptomyces erythreus*). It is a base and readily forms salts with acids. Erythromycin is otherwise known as [(3R\*, 4S\*, 5S\*, 6R\*, 7R\*, 9R\*, 11R\*, 12R\*, 13S\*, 14R\*)-4-[(2,6-dideoxy-3-C-methyl-3-O-methyl- $\alpha$ -L-ribo-hexopyranosyl)-oxy]-  
20 14-ethyl-7,12,13-trihydroxy-3,5,7,9,11,13-hexamethyl-6-[[3,4,6-trideoxy-3-(dimethylamino)- $\beta$ -D-xylo-hexopyranosyl]oxy]oxacyclotetradecane-2,10-dione].

The antibiotic is present in the composition in a pharmaceutically effective concentration. Such concentration  
25 will vary depending on the antibiotic used and the nature of the individual being treated. For guideline purposes, it is recommended that the antibiotic be present at a concentration of about 0.5 to about 6% w/w - composition, preferably about 0.8 to about 5% w/w - composition. In particularly preferred

embodiments of the invention utilizing erythromycin, the erythromycin is at a concentration of about 1.8 to about 4.5% w/w - composition. In preferred embodiments utilizing clindamycin, the clindamycin is at a concentration of about 5 0.7 to about 2% w/w - composition.

In addition to the antibiotic, the package of the present invention can include any other active ingredient that is effective in treating acne and that may be combined with an antibiotic to form a composition which tends to 10 degrade prematurely. Examples of such active ingredients include corticosteroids, tretinoin, antifungal compounds, alpha-hydroxy acids and beta-hydroxy acids.

In preferred embodiments, the acne-treating composition preferably includes an antibiotic as described above and an 15 oxidizing agent which inhibits the formation of free fatty acids in the skin primarily through inactivation of extracellular lipase enzymes. Preferably, the composition includes a "peroxide" oxidizing agent. The term "peroxide" means an organic compound containing an oxygen-oxygen bond 20 capable of cleaving and forming oxygen free-radicals. A variety of peroxides may be used in the practice of the present invention. The peroxides include peroxyacids of carboxylic acids, peroxyesters of carboxylic acids and the dimeric product of carboxylic peroxyacids. Exemplary 25 peroxides include t-butyl peroxyesters of straight and branched chain aliphatic carboxylic acids, and dimeric peroxides such as lauroyl peroxide and benzoyl peroxide.

In preferred embodiments utilizing a peroxide as the oxidizing agent, the peroxide should be of high purity and in

the form of finely divided crystalline particles, preferably micronized particles having a mean average particle size of less than 35 microns. A preferred peroxide for use in the present invention is benzoyl peroxide, and the most preferred  
5 is micronized benzoyl peroxide 70% hydrous.

The oxidizing agent should be present in the composition in a pharmaceutically effective concentration. Such concentration will vary depending on the type of oxidizing agent used and the nature of the individual being treated.  
10 For guideline purposes, it is recommended that the composition contain about 1% to about 30% w/w - composition of oxidizing agent and preferably about 2.5% to about 15% w/w - composition oxidizing agent. In especially preferred embodiments, the oxidizing agent is at a concentration of  
15 about 3 to about 6% w/w - composition.

Active ingredients which are used in formulating the composition of the present invention and which are normally in solid form are advantageously converted into a liquid form in the preparation of the components from which the  
20 composition is made. For example, to prepare a liquid form of an active ingredient, it is combined with an appropriate solvent, the identity of which will depend on the particular active ingredient used. For example, if the active ingredient is highly soluble in water, then water may be used  
25 as the solvent. Water may also be added to the composition as a diluent. If the active ingredient has low solubility in water, another appropriate solvent such as an alcohol may be used as the solvent.

Any pharmaceutically acceptable and suitable material can be used to impart to the component the desired viscosity. Viscosity-enhancing agents are well known. Preferably, a gelling agent is used. The gelling agent may be selected  
5 both as to type and quantity to give products of desired viscosities.

A variety of gelling agents may be used in the practice of the invention. Preferred gelling agents are pure micro-crystalline cellulose, colloidal magnesium-aluminum silicate,  
10 hydroxypropyl methyl cellulose and the so-called hydroxylated vinylic polymers, particularly, those disclosed in U.S. Patent No. 2,798,053. Preferred hydroxylated vinylic polymers are those described generally as interpolymers of a monomeric monolefinic acrylic acid and from about 0.1% to  
15 about 10% by weight based on the total monomer of a monomeric polyether of an oligosaccharide in which the hydroxyl groups that are modified are esterified with allyl groups, with said polyether containing at least two allyl ether groups per oligosaccharide molecule. Commercially available  
20 interpolymers of this type are known as "carbomers" marketed under the trademark Carbopol®. Such interpolymers are described as being polymers of acrylic acid cross-linked with about 1% of a polyallyl ether of sucrose having an average of about 5.8 allyl groups for each sucrose molecule. These  
25 polymers have molecular weight in the order of magnitude of about 1,000,000. Such polymers are available from the B.F. Goodrich Chemical Company and are sold under such trademarks as Carbopol® 934, Carbopol® 940, and Carbopol® 941. A particularly preferred Carbopol is Carbopol® 934 NF.

The gelling agent should be present in the composition at a concentration that provides sufficient viscosity to the composition to allow the composition to adhere to the skin for a sufficient period of time to allow the active  
5 ingredients to act on the affected areas. The effective concentration will depend on the particular gelling agent used and the nature of the component to be gelled. In some embodiments, two or more gelling agents will be present. For guideline purposes, it is recommended that the gelling  
10 agent(s) comprise about 1 to about 4% w/w - composition, preferably about 1.5 to about 3% w/w - composition.

In a highly preferred composition of the present invention, hydroxypropylcellulose is used as a gelling agent.

In a preferred embodiment, a carbomer such as the  
15 Carbopol<sup>®</sup> interpolymers described hereinabove are used as a gelling agent in the oxidizing agent component and hydroxypropylcellulose is used as a gelling agent in the antibiotic component. Hydroxypropylcellulose is a propylene glycol ether of cellulose.

20 The preferred hydroxypropylcellulose used in the practice of the invention is available from Aqualon as Klucel<sup>®</sup> GF having a weight average molecular weight of 370,000.

A surface active agent may be included in the  
25 composition. The surface active agent functions as a processing aid which facilitates even distribution of the oxidizing agent (such as benzoyl peroxide) in the oxidizing agent component gel. The surface active agent may also function to retard the degradative interaction between an

oxidizing agent and an antibiotic. Any pharmaceutically acceptable surface active agent may be used. An example of such a surface active agent is dioctyl sodium sulfosuccinate. The surface active agent is present at a concentration

5 sufficient to facilitate even distribution of the oxidizing agent and optionally to retard the aforementioned degradative interaction. For guideline purposes, it is recommended that the surface active agent comprise about 0.05% to about 1% w/w - composition. In preferred embodiments, dioctyl sodium

10 sulfosuccinate 75% is present at a concentration of about 0.07 to about 0.3% w/w - composition.

The preferred composition includes also a base. The base is added to neutralize the gelling agent (carbomer). The preferred pH range for the carbomer containing gels is about

15 3 to about 7. For guideline purposes, the concentration is about 0.1 to about 1% w/w - composition, preferably about 0.15 to about 0.4% w/w - composition. Suitable bases include, but are not limited to, sodium hydroxide. Other bases, including amines and hydroxides, may be used, for

20 example, triethanolamine, PEG-15 cocamine and potassium hydroxide. In especially preferred embodiments, sodium hydroxide is used at a concentration of about 0.2 to about 0.35% w/w - composition.

Examples of other optional ingredients that may be

25 present in an acne-treating composition are as follows. The composition can include a moistening agent (for example, propylene glycol) at a concentration that functions as a solvent for methyl- and propylparabens. Preferably the moistening agent comprises about 0.5 to about 4% w/w -

composition, most preferably about 1.5 to about 2.5% w/w - composition. The composition may include also an anti-microbial preservative (for example, methylparaben and propylparaben) at a concentration sufficient to inhibit 5 microbial growth in the composition. Preferably, the composition comprises about 0.01 to about 1% w/w - composition, most preferably about 0.03 to about 0.07% w/w - composition.

As mentioned above, a preferred composition for use in 10 the practice of the present invention includes pharmaceutically effective amounts of benzoyl peroxide and erythromycin and hydroxypropylcellulose. A particularly preferred form of this composition comprises: (A) benzoyl peroxide at a concentration of about 2.5 to about 15% w/w - 15 composition, most preferably about 3 to about 6% w/w - composition; (B) erythromycin at a concentration of about 0.8 to about 5% w/w - composition, most preferably about 1.8 to about 4.5% w/w - composition; and (C) hydroxypropylcellulose at a concentration of about 0.5 to about 4% w/w - 20 composition, most preferably about 1 to about 3% w/w - composition.

Preferably the composition includes an additional gelling agent, preferably Carbopol® 934 at a concentration of about 0.5 to about 2% w/w - composition, most preferably 25 about 0.7 to about 1.5% w/w - composition. In this particularly preferred form of the composition, water is used to dissolve the benzoyl peroxide at a concentration of about 35 to about 45% w/w - composition, most preferably about 42 to about 44% w/w - composition and an alcohol, preferably



ethanol (SD #40B Alcohol as described in 27 CFR §21.76) is used to dissolve the erythromycin at a concentration of about 40 to about 48% w/w - composition, most preferably about 43 to about 45% w/w - composition. Optional, but preferred ingredients, of this particularly preferred form of the composition comprise sodium dioctyl sulfosuccinate 75% at a concentration of about 0.05 to about 0.5% w/w - composition, most preferably about 0.08 to about 0.12% w/w - composition and sodium hydroxide at a concentration of about 0.05 to about 0.4% w/w - composition, most preferably about 0.1 to about 0.3% w/w - composition.

There follows a description of exemplary components which can be combined to form the acne-treating composition of the present invention.

Compositions for use in the practice of the present invention are prepared by combining at least two separate active ingredient-containing components. Preferably, a first component used to prepare the composition contains an oxidizing agent and the second component contains an antibiotic. In preferred embodiments of the acne-treating composition, the oxidizing agent is present in equal or greater amounts than the antibiotic, preferably the ratio of oxidizing agent to antibiotic being about 1:1 to about 30:1. In highly preferred embodiments, the ratio of oxidizing agent to antibiotic is about 1:1 to about 5:1.

In preferred embodiments of the present invention, the antibiotic component comprises clindamycin or erythromycin, a solvent for the antibiotic, and a gelling agent. Additional

ingredients which may be present include moistening agents, anti-microbial agents and a base.

In a preferred antibiotic component containing clindamycin ("clindamycin component"), the clindamycin is in the form of clindamycin phosphate and is present in a concentration such that the acne-treating composition includes a pharmaceutically effective concentration of the clindamycin phosphate. A milligram of clindamycin phosphate contains approximately 841 $\mu$ g of clindamycin. Accordingly, a 0.7-2% w/w-composition clindamycin will require approximately 0.8-2.3% w/w-composition clindamycin phosphate. Preferably the clindamycin phosphate concentration comprises about 1.5 to about 3.5% w/w - component. The clindamycin component includes additionally a solvent for the clindamycin. In preferred embodiments, the solvent is purified water which comprises about 88 to about 93% w/w - component. The clindamycin component also includes a gelling agent which imparts thereto a viscosity (about 300,000 to about 500,000 cp) which is similar to the viscosity of the other component used in the composition (preferably a benzoyl peroxide-containing component). The preferred gelling agent is Carbopol® 934 which comprises about 1 to about 2% w/w - component. The clindamycin component can also include a base for the purpose of neutralizing the carbomer (Carbopol® 934) gelling agent. The desired pH range for the carbomer gels is about 3 to about 7. Preferably the base is sodium hydroxide in a concentration of about 0.1 to about 0.3 w/w - component. Other preferred ingredients in the clindamycin component are:

(A) the moistening agent propylene glycol at a concentration

of about 2 to about 6% w/w - component; and (B) the antimicrobial agents methylparaben and propylparaben at concentrations of about 0.04 to about 0.2% w/w - component.

In a preferred antibiotic component containing  
5 erythromycin, the erythromycin is present in a concentration such that the acne-treating composition includes a pharmaceutically effective concentration of the erythromycin. Preferably the erythromycin concentration comprises about 5.5 to about 8.5% w/w - component. As erythromycin has limited  
10 solubility in water, the erythromycin is dissolved in alcohol, preferably SD alcohol #40-B 190 proof, in a concentration of about 87 to about 92% w/w - component. The erythromycin component also includes a gelling agent which imparts to the component a viscosity (about 200,000 cps to  
15 about 400,000 cps) that is similar to the viscosity of the other component used in the composition (preferably a benzoyl peroxide-containing component). Hydroxypropylcellulose is the preferred gelling agent and is present preferably in a concentration of about 3 to about 5% w/w - component.

20 A preferred oxidizing agent component comprises benzoyl peroxide (for example, 70% hydrous) in a concentration such that the acne-treating composition contains a pharmaceutically effective concentration of the benzoyl peroxide. Preferably the benzoyl peroxide comprises about 8  
25 to about 12% w/w - component. The benzoyl peroxide is preferably dissolved in water at a concentration of about 80 to about 85% w/w - component. The oxidizing agent component also includes a gelling agent which imparts to the component a viscosity (about 200,000 to about 500,000 cp) which is

similar to the viscosity of the other component used to prepare the acne-treating composition (preferably an antibiotic component). A preferred gelling agent is Carbopol® 934 at a concentration of about 1 to about 3% w/w - component. The preferred oxidizing agent component comprises also the surface active agent dioctyl sodium sulfosuccinate at a concentration of about 0.1 to 0.3% w/w - component. Another preferred ingredient of the oxidizing agent component is sodium hydroxide which is present at a concentration of about 0.3 to 0.5% w/w - component.

### Examples

The following examples are illustrative of the present invention. The term "USP" as used herein refers to chemicals that conform to the tolerances set forth in the United States Pharmacopeia/National Formulary. (The term "NF" refers to National Formulary standards.) The United States Pharmacopeia/National Formulary is a compendium of pharmaceutical formulations widely used as a standard reference. The term "USP" refers to United States Pharmacopeia standards. The USP/NF is maintained by the United States Pharmacopeial Convention, Inc. The USP24/NF19 was issued January, 2000.

#### Example 1 - Preparation of Benzoyl Peroxide Gel/Erythromycin Gel Package

##### 25      (A) Benzoyl Peroxide Gel

A benzoyl peroxide gel containing the following ingredients was prepared as follows.

	<u>Ingredients</u>	<u>Weight Percent</u>
	Purified Water, USP	82.495
	Carbopol® 934, NF	1.900
	Sodium Hydroxide, NF	0.405
5	Benzoyl Peroxide, USP 70%	15.000
	Diethyl Sodium Sulfosuccinate 75%	0.200

Purified water was added to a tank (hereinafter, the "Phase A tank") and sodium hydroxide pellets were added to the water. The contents of the tank were mixed until the pellets fully  
10 dissolved in the water and the solution cooled to 22-30°C.

Purified water was introduced into a second tank (hereinafter, the "main tank") and the water was cooled to 22-25°C. Carbopol® 934 was dispensed into the main tank and the tank mixed. The contents of the Phase A tank were slowly  
15 added to the contents of the main tank while continuously mixing the contents of the main tank at a temperature of 22-30°C.

Purified water was added to a third tank (hereinafter, the "phase B tank"). The dioctyl sodium sulfosuccinate was  
20 added to the tank, followed by addition of benzoyl peroxide. The contents of the tank were mixed producing a slurry. The Phase B tank was connected to the main tank and the benzoyl peroxide slurry was pumped into the main tank.

The contents of the main tank were mixed under a 10-15"  
25 psi vacuum and transferred to clean and sanitized bulk storage containers. The resultant gel had a viscosity of 360,000 cps. The viscosity was tested using method 1V-101 Brookfield helipath T-F @ 1.5 rpm.

(B) Erythromycin G 1

An erythromycin gel containing the following ingredients was prepared as follows.

	<u>Ingredients</u>	<u>Weight Percent</u>
5	S.D. Alcohol #40-B 190 Proof	89.10
	Erythromycin, USP	6.90
	Hydroxypropylcellulose, NF	4.00

Specially denatured (SD) Alcohol #40-B 190 proof was added to a vessel. Erythromycin was added while mixing.

10 Mixing was continued until the erythromycin completely dissolved. Hydroxypropylcellulose (Klucel® GF manufactured by Aqualon) was then added while mixing. Mixing was performed at -15 to -30 psi to facilitate deaeration. Mixing may be performed in an open or closed vessel. For commercial

15 preparations, mixing will be performed in a closed vessel. After mixing, the tank was sealed and held for a minimum of 12 hours at room temperature in order to allow any entrapped air to dissipate and to assure that all of the hydroxypropylcellulose had gelled. Any evaporated alcohol

20 was replaced with SD Alcohol #40-B to the target weight and the mixture was mixed at -15 to -30 psi. Mixing may be performed in an open or closed vessel. For commercial preparations, mixing will be performed in a closed vessel. The mixture was then transferred into clean and sanitized

25 steel drums. The resultant gel had a viscosity of 360,000 cps. The viscosity was tested using method 1V-101 Brookfield helipath T-F @ 1.5 rpm.

(C) Packaging the Gels

The packaging is performed using a Bartelt horizontal, form-fill-and-seal machine.

The benzoyl peroxide and the erythromycin gels were transferred to pressure vessels charged by inert gas. A pressure regulator was used to keep the pressure level constant. The pressure level was set up to 80 psi for benzoyl peroxide and up to 40 psi for erythromycin gel. A vertical linear piston-metering chamber was installed in each of the gel filling lines. The volume of gel in the chamber was kept constant by the pressurized vessels continuously forcing gel into the metering chamber which assures filling precision and accuracy of viscous gels at high speeds and low volumes. The stroke of the piston in the metering chamber is initiated by a programmable logic controller (PLC) signal synchronized with the movement of the rest of the filling line. The piston drives the gels to a filling pump. A horizontal linear positive displacement micro-pump (*Hi-Bar pump*), with a packet-dosing chamber, was installed at the terminal end of the filling line. The dosing chamber was filled by the action of the metering chamber piston. The dosing chamber feeds a loop of clear tubing terminating in a vertical piston, at the filling station.

The gels are packaged in a two-packet laminated foil package containing the benzoyl peroxide gel in one packet and the erythromycin gel in the other packet. The benzoyl peroxide gel (0.425g) was filled into one packet while the erythromycin gel (0.425g) was simultaneously filled into the second packet. The dispensed gels are within plus or minus

0.025g of the target weight of 0.425g, allowing for a 6% tolerance. After each packet was filled and sealed, the packets were folded and cold-sealed together using a glue adhesive. The packets are constructed of a laminated foil consisting of an inner barrier of low-density polyethylene; a tie layer of ethylene/acrylic acid copolymer; a main barrier of foil; a tie layer of ethylene/acrylic acid copolymer; and an outer barrier of polyester.

Combining the benzoyl peroxide gel with the erythromycin gel provides an acne-treating composition having the following ingredients by weight%.

INGREDIENT	% W/W COMPOSITION
Purified Water, USP	43.50
Carbopol® 934, NF	0.95
Sodium Hydroxide, NF	0.20
Benzoyl Peroxide, 70% Hydrous, USP	5.25^
Dioctyl Sodium Sulfosuccinate, DF	0.10
SD Alcohol #40-B/190°, DF	44.55
Erythromycin, USP	3.45
Hydroxypropylcellulose, NF	2.00

**Example 2 - Preparation of Benzoyl Peroxide/Clindamycin Gel Package**

The benzoyl peroxide gel is prepared as described in Example 1.

A clindamycin gel containing the following ingredients was prepared as follows.



	<u>Ingredients</u>	<u>Weight Percent</u>
	Purified Water, USP	91.28
	Carbomer 934	1.50
	Sodium Hydroxide, NF	0.27
5	Propylene Glycol, USP	4.00
	Methylparaben, NF	0.14
	Propylparaben, NF	0.06
	Clindamycin Phosphate, USP	2.75

Purified water was added to a tank (hereinafter, the  
10 "Phase A tank") and sodium hydroxide pellets were added to  
the purified water and the contents of the tank were mixed  
until the pellets fully dissolved in the purified water and  
the solution cooled to 22-30°C.

Purified water was added into a second tank  
15 (hereinafter, the "main tank") and Carbomer 934 was dispensed  
into the main tank. The contents of the phase A tank were  
slowly added to the contents of the main tank while  
continuously mixing the contents of the main tank and  
deaerating under a vacuum. The temperature was maintained at  
20 22-30°C.

Purified water was added to a third tank (hereinafter,  
the "phase B tank"). Clindamycin phosphate was added and  
mixed until dissolved. Once the clindamycin phosphate was  
completely dissolved, the contents of the tank were pumped  
25 into the main tank and mixed under vacuum.

Purified water and propylene glycol were added to a  
fourth tank. The tank was heated to 35-40°C. Methylparaben  
was then added and the contents of the tank were mixed until  
all the methylparaben was dissolved. Propylparaben was then  
30 added and the contents of the tank mixed until all the  
propylparaben was dissolved. The contents were transferred  
into the main tank.

While continuously mixing, a vacuum was turned on and the mixing continued until the material was homogenous. The contents of the tank were then transferred to stainless steel drums. The resultant gel had a viscosity of 370,000 cps.

- 5 The viscosity was tested using method IV-101 Brookfield helipath T-P @ 1.5 rpm.

The benzoyl peroxide and the erythromycin gels were transferred to pressure vessels charged by inert gas. A pressure regulator was used to keep the pressure level  
10 constant. The pressure level was set up to 80 psi for benzoyl peroxide and up to 40 psi for erythromycin gel. A vertical linear piston-metering chamber was installed in each of the gel filling lines. The volume of gel in the chamber was kept constant by the pressurized vessels continuously  
15 forcing gel into the metering chamber which assures filling precision and accuracy of viscous gels at high speeds and low volumes. The stroke of the piston in the metering chamber is initiated by a PLC signal synchronized with the movement of the rest of the filling line. The piston drives the gels to  
20 a filling pump. A horizontal linear positive-displacement micro-pump, with a packet dosing chamber, was installed at the terminal end of the filling line. The dosing chamber was filled by the action of the metering chamber piston. The dosing chamber feed a loop of clear tubing terminating in a  
25 vertical piston, at the filling station.

For the final packaging step, 0.425g of a 10% benzoyl peroxide gel was filled into one packet while 0.425g of a 2% clindamycin gel was simultaneously filled into the second packet. The dispensed gels are within plus or minus 0.025 g

of the target weight of 0.425g, allowing for a 6% tolerance. After each pouch was filled and sealed, the pouches were folded and cold-sealed together.

Combining the benzoyl peroxide gel with the clindamycin gel provides an acne-treating composition having the following ingredients by weight%.

	INGREDIENT	% W/W COMPOSITION
	Purified Water, USP	89.14
	Carbomer 934, NF	1.70
10	Sodium Hydroxide, NF	0.34
	Propylene Glycol, USP	2.00
	Diethyl Sodium Sulfosuccinate 75%, DF	0.10
	Methylparaben, NF	0.07
	Propylparaben, NF	0.03
15	*Clindamycin Phosphate, USP	1.37
	Hydrous Benzoyl Peroxide, USP	5.25

\*The % w/w for the clindamycin phosphate is based solely on % w/w. The active substance in the product is clindamycin. Clindamycin phosphate is converted *in vitro* to active clindamycin. The actual amount of clindamycin as clindamycin phosphate is based on several factors including activity. The actual amount of material added is based on 6.875 kg of clindamycin phosphate.

Claims

1. A package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, said composition tending to degrade prematurely, one of said components comprising an oxidizing agent and another of said components comprising an antibiotic which is effective against acne-associated bacterial species, said components separated one from the other in said package, said one component having a viscosity within about 50% of the viscosity of said other component.

2. The package of Claim 1 wherein said component comprising an oxidizing agent comprises about 8 to about 12% w/w - component oxidizing agent.

3. The package of Claim 1 wherein said component comprising an antibiotic comprises about 1.5 to about 8.5% w/w - component antibiotic.

4. The package of Claim 1 wherein the ratio of said oxidizing agent to said antibiotic is about 1:1 to about 30:1.

5. The package of Claim 4 wherein the ratio of said oxidizing agent to said antibiotic is about 1:1 to about 5:1.

6. The package of Claim 1 wherein said oxidizing agent is benzoyl peroxide.

7. The package of Claim 1 wherein said antibiotic is selected from the group consisting of erythromycin and clindamycin.

8. The package of Claim 7 wherein said antibiotic is erythromycin.

9. The package of Claim 1 wherein said package comprises a dual packet laminated foil package.

10. A package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of said components comprising a benzoyl peroxide gel and another of said components comprising a gel of erythromycin, said components separated one from the other, said benzoyl peroxide gel having a viscosity within about 50% of the viscosity of said erythromycin gel.

11. The package of Claim 10 wherein said benzoyl peroxide gel comprises about 8 to about 12% w/w - component benzoyl peroxide.

12. The package of Claim 11 wherein said benzoyl peroxide gel comprises about 10% w/w - component benzoyl peroxide.

13. The package of Claim 10 wherein said erythromycin gel comprises about 5.5 to about 8.5% w/w - component erythromycin.

14. The package of Claim 13 wherein said erythromycin gel comprises about 7% w/w - component erythromycin.

15. The package of Claim 14 wherein the ratio of said benzoyl peroxide to said erythromycin is about 1:1 to about 30:1.

16. The package of Claim 15 wherein the ratio of said benzoyl peroxide to said erythromycin is about 1:1 to about 5:1.

17. The package of Claim 10 wherein said package is a dual packet laminated foil package.

18. A package comprising: (A) components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of said components comprising a benzoyl peroxide gel and another of said components comprising a gel of erythromycin and hydroxypropylcellulose; and (B) containers for holding said components in the package separated one from the other; said components having viscosities such that, upon the application of a uniform force to said components, substantially equal volumes of said components are capable of being dispensed simultaneously from said containers.

19. The package of Claim 18 wherein said benzoyl peroxide gel comprises about 8 to about 12% w/w - component benzoyl peroxide.

20. The package of Claim 18 wherein said erythromycin and hydroxypropylcellulose gel comprises about 5.5 to about 8.5% w/w - component erythromycin and about 3 to about 5% w/w - component hydroxypropylcellulose.

21. The package of Claim 18 wherein said benzoyl peroxide gel comprises about 10.5% w/w - component benzoyl peroxide and said gel of erythromycin and hydroxypropylcellulose comprises about 7% w/w - component erythromycin and about 4% w/w - component hydroxypropylcellulose.

22. The package of Claim 21 wherein said benzoyl peroxide gel has a viscosity from about 200,000 cps to about 500,000 cps and said erythromycin and hydroxypropylcellulose gel has a viscosity of from about 200,000 cps to about 400,000 cps.

23. A composition for the treatment of acne comprising:

- (a) benzoyl peroxide;
- (b) erythromycin; and
- (c) hydroxypropylcellulose.

24. The composition of Claim 23 including an additional gelling agent.

25. The composition of Claim 23 including a surface active agent.

26. The composition of Claim 23 including a base which increases the pH of said composition to a value at which said composition has an acceptable viscosity.

27. The composition of Claim 23 including an alcohol solvent for said erythromycin.

28. The composition of Claim 23 including a diluent.

29. A composition for the treatment of acne comprising: (a) benzoyl peroxide; (b) erythromycin; (c) hydroxypropylcellulose; (d) a gelling agent; (e) a surface active agent; (f) a base; (g) an alcohol; and (h) water.

30. The composition of Claim 29 wherein said gelling agent is a hydroxylated vinylic polymer.

31. The composition of Claim 29 wherein said surface active agent is dioctyl sodium sulfosuccinate.

32. The composition of Claim 29 wherein said base is sodium hydroxide.

33. The composition of Claim 29 wherein said alcohol is specially denatured ethanol.



34. A composition for the treatment of acne comprising about 5.25% w/w - composition benzoyl peroxide, about 3.45% w/w - composition erythromycin, and about 2% w/w - composition hydroxypropylcellulose.

35. The composition of Claim 34 comprising about 0.95% w/w - composition of a hydroxylated vinylic polymer, about 0.10% w/w - composition of dioctyl sodium sulfosuccinate, about 0.2% w/w - composition of sodium hydroxide, about 44.55% w/w - composition specially denatured ethanol and about 43.5% w/w - composition of water.


36. A package containing a first acne-treating composition and a second acne-treating composition, said compositions tending to degrade chemically upon being mixed with one another, said package comprising a first chamber containing said first composition and a second chamber containing said second composition, each of said chambers having a sealed dispensing orifice, said orifices positioned adjacent to one another.

37. A package containing a first acne-treating composition and a second acne-treating composition, said compositions tending to degrade chemically upon being mixed with one another, said package comprising a first and second packet, said packets being attached to each other and each of said packets having a sealed dispensing orifice, said sealed dispensing orifices being positioned adjacent to each other


so that, as said compositions are dispensed through the orifices in unsealed form, they are combined with each other.


38. A package containing a first acne-treating composition and a second acne-treating composition, said compositions tending to degrade chemically upon being mixed with one another, said package comprising a first and second packet, each of said packets having a top which includes a dispensing orifice that has a tear-off tab seal, a bottom, two sides in a parallel relationship to one another, and an exterior face and an interior face, said first packet containing said first composition and said second packet containing said second composition, said interior faces of said packets adjacent to and facing one another such that said tear-off tabs can be simultaneously torn off such that said compositions are dispensed and combined with each other as they exit said packets through said dispensing orifices.

**FIGURE 1**



**FIGURE 2**



**FIGURE 3**





FIGURE 4

**FIGURE 5**

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/JS 01/18033

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 A61K9/06 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 088 717 A (RORER INTERNATIONAL) 16 June 1982 (1982-06-16) claims 1-8, 11, 16-20	1-19, 36-38
Y	page 3, line 27 - line 47 examples 3-6	20-35
Y	US 4 954 487 A (EUGENE R. COOPER; ET AL.) 4 September 1990 (1990-09-04) examples 17, 18	20-35
X	WO 99 02133 A (GIST BROCADES) 21 January 1999 (1999-01-21) claims 1-6 page 3, line 17 - line 24 page 4, line 17 - line 21 page 4, line 34 - line 37 page 6, line 9 - line 21 example 1	1-19, 36-38

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

13 November 2001

Date of mailing of the international search report

21/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JS 01/18033

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 93 15726 A (BAROODY, LLOYD; ET AL.) - 19 August 1993 (1993-08-19) claims 1,2 examples 1-13 ----	1-19, 36-38
X	CA 2 015 111 A (WOO, TOM; ET AL.) 23 October 1991 (1991-10-23) claims 1,4,5,11,13,72-79 -----	1-19, 36-38



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/18033

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2088717	A	16-06-1982	AR 229405 A1	15-08-1983
			AT 381452 B	27-10-1986
			AT 285581 A	15-03-1986
			AU 557147 B2	11-12-1986
			AU 7202981 A	17-06-1982
			BE 889327 A1	16-10-1981
			CA 1179599 A1	18-12-1984
			CH 656307 A5	30-06-1986
			CH 647147 A5	15-01-1985
			DE 3124698 A1	08-07-1982
			DK 273981 A	09-09-1982
			ES 503797 D0	01-03-1983
			ES 8303087 A1	01-05-1983
			FR 2495471 A1	11-06-1982
			GB 2090135 A	07-07-1982
			IE 52227 B1	19-08-1987
			IT 1210608 B	14-09-1989
			JP 57099525 A	21-06-1982
			MX 9203750 A1	01-09-1992
			MX 6555 E	10-07-1985
			NL 8102997 A	01-07-1982
			NZ 197491 A	28-02-1985
			NZ 209389 A	28-02-1985
			PH 19245 A	14-02-1986
			SE 8103925 A	09-06-1982
			US 4497794 A	05-02-1985
			US 4692329 A	08-09-1987
			ZA 8104215 A	23-02-1983
US 4954487	A	04-09-1990	EP 0013459 A2	23-07-1980
			JP 55120514 A	17-09-1980
			AU 544969 B2	27-06-1985
			AU 7272081 A	14-01-1982
			CA 1165240 A1	10-04-1984
			DE 3172516 D1	07-11-1985
			EP 0043738 A2	13-01-1982
			IE 51377 B1	10-12-1986
			JP 1737953 C	26-02-1993
			JP 4020886 B	07-04-1992
			JP 57081408 A	21-05-1982
			NZ 197668 A	16-03-1984
			ZA 8104650 A	28-07-1982
WO 9902133	A	21-01-1999	AU 9259598 A	08-02-1999
			BR 9810985 A	15-08-2000
			CN 1262616 T	09-08-2000
			WO 9902133 A2	21-01-1999
			EP 0996428 A2	03-05-2000
			JP 2001509477 T	24-07-2001
WO 9315726	A	19-08-1993	AT 197248 T	15-11-2000
			AU 3478893 A	03-09-1993
			BR 9305884 A	19-08-1997
			CA 2130341 A1	19-08-1993
			CN 1077887 A ,B	03-11-1993
			DE 69329617 D1	07-12-2000
			DE 69329617 T2	23-05-2001
			DK 626844 T3	22-01-2001

## INTERNATIONAL SEARCH REPORT

International Application No

PCT 01/18033

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 9315726	A	EP 0626844 A1	07-12-1994	
		EP 0960618 A2	01-12-1999	
		ES 2153377 T3	01-03-2001	
		FI 943783 A	17-08-1994	
		JP 7504168 T	11-05-1995	
		MX 9300848 A1	31-08-1994	
		NO 943045 A	17-08-1994	
		NZ 249021 A	27-08-1996	
		PL 171127 B1	28-03-1997	
		PT 626844 T	30-03-2001	
		RU 2122412 C1	27-11-1998	
		SG 48195 A1	17-04-1998	
		WO 9315726 A1	19-08-1993	
		US 5733886 A	31-03-1998	
		US 6117843 A	12-09-2000	
CA 2015111	A	23-10-1991	CA 2015111 A1	23-10-1991

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
**WO 01/091726 A1**(51) International Patent Classification<sup>7</sup>: A61K 9/06, 7/48

(21) International Application Number: PCT/US01/18033

(22) International Filing Date: 1 June 2001 (01.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/209,330 2 June 2000 (02.06.2000) US  
0023891.5 29 September 2000 (29.09.2000) GB(71) Applicant (for all designated States except US): **DERMIK INTERNATIONAL HOLDING INC.** [US/US];  
3711 Kennett Pike, Suite 200, Greenville, DE 19807 (US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): **MANETTA, Vincent, E.** [US/US]; Galderma, 5 Cedarbrook Drive, Cranbury, NJ 08512 (US). **MAHONEY, Linda, M.** [US/US]; 103 Beaufort Court, Phoenixville, PA 19460 (US). **NORDSIEK, Michael, T.** [US/US]; 561 Bair Road, Berwyn, PA 19312 (US). **BALAJI, Kodumudi, S.** [US/US]; 1521 Ide Drive, Lansdale, PA 19446 (US). **DORRELL, Jay** [US/US]; 1459 Woodford Drive, Wayne, PA 19087 (US).(74) Agents: **BARRON, Alexis et al.**; Synnestvedt & Lechner LLP, 2600 Aramark Tower, 1101 Market Street, Philadelphia, PA 19107-2950 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

[Continued on next page]

(54) Title: ACNE-TREATING COMPOSITION





CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

**Declaration under Rule 4.17:**

— *of inventorship (Rule 4.17(iv)) for US only*

**Published:**

— *with international search report*

**(48) Date of publication of this corrected version:**

15 August 2002

**(84) Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**(15) Information about Correction:**

see PCT Gazette No. 33/2002 of 15 August 2002, Section II

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

-1-

**ACNE-TREATING COMPOSITION****Field of the Invention**

The present invention relates to a pharmaceutical composition that tends to lose its effectiveness prematurely  
5 and to improved means for the effective use of such a composition. More particularly, the present invention relates to the packaging of constituents comprising such a composition and to an improved form of such composition, including, for example, a composition for topical  
10 application, such as a topical composition for the treatment of acne.

Acne is a common inflammatory disease which is very common at puberty and may continue for many years. It occurs in facial skin areas where sebaceous glands are the  
15 largest, most numerous, and most active. In its milder forms, acne is a superficial disorder which is evidenced by slight, spotty irritations and which can be treated satisfactorily by ordinary skin hygiene. However, in the more inflammatory types of acne, bacterial invasion of or  
20 about the pilosebaceous follicles occurs and results in

- 2 -

the formation of pustules, infected cysts, and, in extreme cases, canalizing inflamed and infected sacs appear.

These lesions may become extensive and leave permanent, disfiguring scars.

5           Therapeutic methods for treating acne are designed to prevent formation of new lesions and facilitate the healing of old lesions. Treatments include the systemic and topical administration of anti-acne agents such as antibiotics or synthetic Vitamin A analogs. In most  
10 cases, systemic treatment of acne is not desirable because of the risks of adverse side effects that are experienced by the user. For this reason, topical acne treatment compositions have been preferred.

The present invention relates to topical anti-acne  
15 compositions and packages which contain the components of the composition.

### Reported Developments

Topical anti-acne compositions include, for example, one or more of the following: sulfur, resorcinol,  
20 salicylic acid, benzoyl peroxide, steroids and antibiotics. Exemplary antibiotics are disclosed in U.S. Patent No. 3,969,516 (lincomycin family); BR Publication No. 1,594,314 (erythromycin); and U.S. Patent No. 3,952,099 (tetracycline). Preparations containing a  
25 peroxide are reported in U.S. Patent Nos. 3,535,422; 4,056,611; 4,387,107, and British Publication No. 1,594,314 and U.S. Patent No. 4,497,794. Antibiotic-

- 3 -

containing compositions which also include anti-inflammatory steroids are disclosed in U.S. Patent No. 4,132,781.

Attempts to improve the effectiveness of topical  
5 antibiotic compositions for use in the treatment of acne have taken a number of approaches. One approach is reported in U.S. Patent Nos. 3,989,815; 3,989,816, 3,991,203; 4,122,170; 4,316,893; 4,444,762; and EP 27,286, which disclose skin-penetrating vehicle compositions that  
10 reportedly increase the transdermal absorption of any physiologically active substance, including antibiotics. However, some penetrating agents are not compatible with antibiotics used to treat acne.

A further approach relates to the use of a  
15 composition which utilizes two different active components, such as erythromycin, clindamycin, Vitamin A acid or benzoyl peroxide. Compositions including mixtures of a peroxide and erythromycin are reported in British Publication No. 1,594,314. Acne-treating compositions  
20 comprising benzoyl peroxide, erythromycin, and hydroxyethylcellulose have also been disclosed in "The Formulation and Stability of Erythromycin-Benzoyl Peroxide in a Topical Gel, Vermeulen et al., *International Journal of Pharmaceutics*, 178, 137-141 (1999)". U.S. Patent No.  
25 5,894,019 to Hess et al. discloses a topically applied pharmaceutical composition that is prepared by mixing a liquid solution of benzoyl peroxide (or other pharmaceutically active ingredient) that is dissolved in a

- 4 -

liquid lipid with a hydrous gel that may also contain a pharmaceutically active ingredient, such as erythromycin. The gelling agent used to form the hydrous gel may include hydroxypropylcellulose. U.S. Patents Nos. 5,466,446 to 5 Stiefel et al, 5,733,886 to Barody et al., and 5,767,098 to Klein et al. disclose acne-treating compositions which contain clindamycin and benzoyl peroxide.

The above compositions comprising a peroxide, such as benzoyl peroxide, and an antibiotic, such as erythromycin 10 or clindamycin, may have stability limitations. When the peroxide and antibiotic are combined, the peroxide can partially or completely oxidize the antibiotic, rendering it ineffective. If the peroxide and antibiotic are provided separately, it is usually necessary to have a 15 pharmacist combine the component containing the peroxide with the component containing the antibiotic to ensure that the composition is prepared properly, has the correct final concentrations of the peroxide and antibiotic, and to maximize the shelf life of the composition by 20 postponing the mixing of the components until the time of sale. Another problem with these compositions is that once mixed, the oxidation reactions of the peroxide not only degrade the antibiotic, but also result in the accumulation of undesirable degradative products in the 25 composition. An additional problem is that unused portions of certain acne-treating composition must be refrigerated to minimize degradation until use.



- 5 -

The present invention relates to improved means for using effectively anti-acne compositions that tend to lose their effectiveness prematurely and to improved forms of anti-acne compositions.

5                    Summary of the Invention

In accordance with the present invention, there is provided a package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, the composition  
10    tending to degrade prematurely, one of the components comprising an oxidizing agent and another of the components comprising an antibiotic which is effective against acne-associated bacterial species, the components separated one from the other in the package, one component  
15    having a viscosity within about 50% of the viscosity of the other component.                    In preferred embodiments, the oxidizing agent is benzoyl peroxide, the antibiotic is selected from the group consisting of erythromycin and clindamycin, and the package is a dual  
20    packet laminated foil package.

Another aspect of the present invention provides a package comprising: (A) components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of the components  
25    comprising a benzoyl peroxide gel and another of the components comprising a gel of erythromycin and hydroxypropylcellulose; and (B) containers for holding the

- 6 -

components in the package separated one from the other; the components having viscosities such that, upon the application of a uniform force to the components, substantially equal volumes of the components are capable  
5 of being dispensed simultaneously from the containers.

Yet another aspect of the present invention is a composition for the treatment of acne comprising: (A) benzoyl peroxide; (B) erythromycin; and (C) hydroxypropylcellulose. Examples of preferred  
10 compositions may further comprise an additional gelling agent, an agent which inhibits degradative interactions between benzoyl peroxide and erythromycin, a base which increases the pH of the composition to a value at which said composition has an acceptable viscosity, an alcohol  
15 solvent for erythromycin, and a diluent.

Yet another aspect of the present invention provides a composition for the treatment of acne comprising: (a) benzoyl peroxide; (b) erythromycin; (c) hydroxypropylcellulose; (d) a gelling agent; (e) a surface  
20 active agent; (f) a base; (g) an alcohol; and (h) water. An example of a preferred gelling agent includes a hydroxylated vinylic polymer.

An example of a preferred base is sodium hydroxide. An example of a preferred alcohol used in the composition  
25 is specially denatured ethanol. An example of a surface active agent is dioctyl sodium sulfosuccinate.

The invention includes also within its scope the provision of a package containing a first acne-treating

- 7 -

composition and a second acne-treating composition, the compositions tending to degrade chemically upon being mixed with one another, the package comprising a first chamber containing the first composition and a second  
5 chamber containing the second composition, each of the chambers having a sealed dispensing orifice, the orifices positioned adjacent to one another.

Another aspect of the present invention provides a package containing a first acne-treating composition and a  
10 second acne-treating composition, the compositions tending to degrade chemically upon being mixed with one another, the package comprising a first and second packet, the packets being attached to each other and each of the packets having a sealed dispensing orifice, the sealed  
15 dispensing orifices being positioned adjacent to each other so that, as the compositions are dispensed through the orifices in unsealed form, they are combined with each other.

An additional aspect of the present invention  
20 provides a package containing a first acne-treating composition and a second acne-treating composition, the compositions tending to degrade chemically upon being mixed with one another, the package comprising a first and second packet, each of the packets having a top which  
25 includes a dispensing orifice that has a tear-off tab seal, a bottom, two sides in a parallel relationship to one another, and an exterior face and an interior face,

- 8 -

the first packet containing the first composition and the second packet containing the second composition, the interior faces of said packets adjacent to and facing one another such that the tear-off tabs can be simultaneously  
5 torn off such that the compositions are dispensed and combined with each other as they exit the packets through the dispensing orifices.

### Description of the Drawings

Figure 1 is a plan view of a dual-packet package.

10 Figure 2 is a plan view of the package of Figure 1 folded about axis A-A.

Figure 3 is a cross-sectional view of a trilaminate package.

Figure 4 is an exploded view of a pump for dispensing  
15 the compositions of the present invention.

Figure 5 is a partial sectional view of the assembled pump depicted in Figure 4.

### Detailed Description of the Invention

The acne-treating composition of the present  
20 invention comprises a plurality of components which when combined tend to degrade prematurely. Degradation is considered premature when the components are mixed together and at least one of the components degrades at a rate faster than the rate at which it degrades (if at all)  
25 in the absence of the other component. As an example, in acne-treating compositions which contain an oxidizing

- 9 -

agent and an antibiotic, the oxidizing agent will oxidize and thereby degrade the antibiotic at a much faster rate than the antibiotic would degrade in the absence of the oxidizing agent.

5           Components that degrade prematurely when combined may be used in the practice of the present invention and may be identified using techniques known in the art. Examples of such components include, but are not limited to, oxidizing agents, gelling agents, and antibiotics.

10           The acne-treating composition of the present invention is provided in a package which maintains the components separated from each other until the components are removed in whole or in part from the package. In preferred form, the package is designed to enable the user  
15 to extract readily from the package amounts of components which form an effective acne-treating composition.

          Use of such a package allows long-term storage of the components at room temperature without degradation. The package can be carried conveniently by the patient because  
20 the contents do not require refrigeration. This increases patient compliance with administration of the composition. In addition, because the composition is mixed at the time of application, degradation is minimized. Furthermore, the proportions of active ingredients in the composition  
25 can be optimized by exact filling of packets and by the ease with which the patient can empty the packets simultaneously such that the final composition has the desired ratios of components.

- 10 -

The package of the present invention may be made from any suitable material, for example, paper, plastic, or metal foil, in laminated or composite form. The material functions to protect the components from exposure to air,  
5 moisture, light and other environmental factors which could change the physical and/or chemical nature of the components and to keep the components from contact with each other.

In preferred form, the package of the present  
10 invention includes chambers which maintain the components of the composition in separated form and which accommodate amounts of the components such that, when the components are removed from their chambers and brought together, they form a single effective acne-treating dose of the  
15 composition. For example, in preferred embodiments, the package contains a sufficient amount of the composition to apply a thin layer of the composition to the entire surface of the face (with the exception of the eyes and mouth) and sufficient amounts of the pharmaceutically  
20 active ingredients to treat effectively any acne present in the areas to which the composition is applied.

It is believed that the present invention will be used widely in the packaging of an acne-treating composition comprising a component which includes an  
25 oxidizing agent and a component which includes an antibiotic which is effective in treating an acne condition. As mentioned above, the oxidizing agent in such a composition tends to oxidize and thereby degrade

- 11 -

the antibiotic when the two components are brought together. The present invention may also be used in acne-treating compositions which include retinoic acid.

A particularly preferred embodiment of the package of the present invention is shown in Figure 1, in which there is illustrated a dual-packet, laminated metal foil package 1 comprising a first packet 5 which includes a first chamber 2 containing a first component of the pharmaceutical composition and a second packet 7 which includes a second chamber 3 containing a second component of the composition. Each of the chambers 2 and 3 has a sealed dispensing orifice 9 and 9' with tear-off tabs 11 and 11'. Each packet has a top portion 13 and 13', a bottom portion 15 and 15', a commonly shared side 17 and a second side 19 and 19' in parallel relationship to each other, and an interior face 21 and 21' and an exterior face (not shown in Figure 1) in opposed relationship to each other. The first chamber 2 and second chamber 3 may be of equal or different size.

Figure 2 shows the package of Figure 1 in a form in which the packets have been folded about axis A and the interior faces 21 and 21' of the packet are in apposition to one another and the exterior faces 23 and 23' (not shown) face outwardly. The tear-off tabs 11 and 11' of the packets are superimposed upon one another. This permits them to be removed simultaneously. After removal of the tabs, the components in chambers 2 and 3 can be

- 12 -

dispensed and combined with each other as they exit packets 5 and 7 through the dispensing orifices 9 and 9'.

In preferred embodiments, the packages are comprised of two packets made of a multi-layered laminated foil.

5 Preferably the laminate is a five-layer design consisting of three main barrier layers and two adhesive layers which lie between the barrier layers and hold the layers together. In preferred embodiments, the packets have a first innermost layer comprised of low density  
10 polyethylene. This layer is in contact with the composition. The second layer is an adhesive layer comprised of ethylene-acrylic acid copolymer which binds the first layer to the third layer. The third layer is a barrier layer made of aluminum foil. The fourth layer is  
15 comprised of ethylene-acrylic acid copolymer which serves as an adhesive layer binding the third layer to the fifth layer. The fifth and final layer is comprised of polyester which provides durability and tear- and puncture-resistance to the package. A preferred laminate  
20 for use in the present invention is available from Reynolds Metals Co.

For exemplary purposes, the use of the package of Figures 1 and 2 is described in connection with an acne-treating composition comprising an oxidizing agent and an  
25 antibiotic. Accordingly, one of the chambers of the package contains the oxidizing agent and the other chamber contains the antibiotic. To extract the oxidizing agent and antibiotic from the package, both packets thereof are



- 13 -

opened simultaneously by tearing off their respective tear-off tabs. The packets are then squeezed by the user until the contents of the chambers are emptied substantially. The contents typically exit their  
5 respective orifices in the form of a ribbon. The oxidizing agent and the antibiotic component can be mixed conveniently to form the acne-treating composition by the user's dispensing the contents into the palm of one hand and using the fingers of the other hand to mix the  
10 components together by use of a swirling motion. The composition is then applied to the site or sites in need of treatment.

The package of Figures 1 and 2 can be characterized as a single-dose package which contains a sufficient  
15 amount of each component to form a volume of composition which can be applied as a thin layer to the entire surface of the face (with the exception of the eyes and mouth) with a minimal amount of the composition left over. Any remaining composition may be discarded. Because of the  
20 relatively small amounts of components in the package, the package can be considered also to be a "single-use" package which is thrown away after its contents are emptied.

Figure 3 illustrates a cross-sectional view of  
25 another package embodiment in which there is illustrated a "trilaminate" package 26 comprising a first packet 27 and a second packet 29, having a common internal barrier layer 31, preferably a single layer of aluminum foil or other

- 14 -

suitable material known in the art. This internal barrier layer 31 is in contact with the first 32 and second 33 compositions within the first 27 and second 29 packets.

The internal barrier layer 31 is sealed along its

5 perimeter to a first exterior barrier layer 34, and a second exterior barrier layer 36. Both the first 34 and second 36 exterior barrier layers are preferably comprised of a single layer, of aluminum foil or other suitable material known in the art.

10 A "multiple use" package is capable of holding relatively large amounts of components for use in forming the treating composition. The multiple-use package is similar to the single-use package in that the components of the acne-treating composition are kept separate from  
15 one another and mixed when the product is dispensed. In preferred embodiments, the multiple-use package may include a dose-metering pump which dispenses an equal amount of both components of the acne-treatment composition.

20 Figure 4 shows an exploded side view of a pump 50 for dispensing a multi-component fluid formulation, for example, a topical anti-acne composition. The preferred pump embodiment shown is designed to accommodate a two-component formulation, although it is understood that the  
25 design is not limited to use with formulations having only two components.

Pump 50 is suitable for use with lotions, creams, oils, gels and liquids and is designed to keep the

components of the formulation separate while they are stored in the pump. Mixing of the components occurs only after they have been dispensed from the pump immediately before use. This feature is useful to extend the effective life of formulations, such as the anti-acne composition, which tend to degrade once the components are combined. The pump is also designed to dispense the components of the formulation in the proper relative amounts which provide maximum effectiveness of the formulation.

As seen in Figure 4, pump 50 comprises a bottle 52 having two reservoirs 54a and 54b separated from one another by a bulkhead 56. The reservoirs hold the different components of the formulation and are attached to a common neck 58 which provides for attachment of the other pump components, described below, to the bottle 52. Neck 58 surrounds and defines two openings 60a and 60b in reservoirs 54a and 54b respectively, the openings permitting the components of the formulation to be loaded into the reservoirs. A liner 62 engages neck 58 and seals openings 60a and 60b to prevent premature mixing of the components within the pump.

Liner 62 has two through openings 64a and 64b through which dip tubes 66a and 66b extend into reservoirs 54a and 54b respectively. Dip tubes 66a and 66b provide conduits for drawing the formulation components from the reservoirs 54a and 54b to be dispensed by the pump. Each dip tube is connected to a respective valve seat 68a or 68b, the valve

- 16 -

seats being attached to a receptacle 70 which sits atop neck 58.

As best seen in Figure 5, receptacle 70 has two cylinders 72a and 72b which are in fluid communication with dip tubes 66a and 66b through valve seats 68a and 68b respectively. Housed in each cylinder is a respective ball 74a or 74b. Balls 74a and 74b are preferably stainless steel and are sized to seat in respective valve seats 68a and 68b to form a pair of check valves 76a and 76b which allow components of the formulation to flow in one direction only, from the respective reservoirs 54a and 54b through the dip tubes 66a and 66b and into cylinders 72a and 72b as described in detail below.

Cylinders 72a and 72b each receive a respective piston 78a and 78b. The pistons are sized to sealingly interfit slidably within their respective cylinders and provide suction and pumping action when they are reciprocated within the cylinders during pump actuation. Pistons 78a and 78b have respective piston bores 80a and 80b which provide fluid communication between their respective cylinders 72a and 72b and a spout 82 attached to the tops of the pistons. Pistons 78a and 78b are each biased to a position away from check valves 76a and 76b by respective springs 84a and 84b housed within cylinders 72a and 72b between the valve seats 68a and 68b and the pistons 78a and 78b as shown.

As seen in Figure 5, spout 82 has ducts 86a and 86b which are in fluid communication with piston bores 80a and

- 17 -

80b of pistons 78a and 78b respectively. A second set of valve seats 88a and 88b are provided at the tops of piston bores 80a and 80b. A second set of balls 90a and 90b are positioned within ducts 86a and 86b respectively, the  
5 balls being sized to seat on respective valve seats 88a and 88b and form a second set of check valves 92a and 92b. Preferably, balls 90a and 90b are made of stainless steel. Ducts 86a and 86b are in fluid communication with respective dispensing nozzles 94a and 94b from which the  
10 components held in reservoirs 54a and 54b are dispensed to be mixed outside of pump 50.

As seen in Figures 4 and 5, most of the pump components including the spout 82, pistons 78a and 78b, check valves 76a, 76b, 92a and 92b, springs 84a and 84b,  
15 and receptacle 70 are housed within a retainer 96. Retainer 96 has a lower portion 98 adapted to engage neck 58 and retain the pump components to the bottle 52. Attachment of the retainer is preferably by an interference fit between the retainer inner surface 100  
20 and raised ribs 102 surrounding neck 58. An over cap 104 fits over the retainer 96 and protects the moving parts of the pump against unintentional pump actuation.

Pump 50 is preferably made from various types of resilient, flexible plastic materials, such as  
25 polyethylene and polypropylene to cite examples. Plastics are advantageous because they are inexpensive, easily molded, substantially unbreakable, and chemically inert and, thus, will not affect the components of the

- 18 -

formulation. Parts of the pump, such as the springs for biasing the pistons and the balls for the check valves, for which plastic is not suited are preferably made of stainless steel. Stainless steel provides a robust, durable material which also is substantially chemically inert and, thus, suitable for pump parts which will come into contact with the components of the formulation during pump operation.

Operation of the dual dispensing pump 50 is described with reference to Figure 5. In operation, the user removes the over cap 104 exposing spout 82. With the pump 50 in an upright position, the user manually depresses spout 82. This causes the spout and attached pistons 78a and 78b to move downwardly with respect to the retainer 96, check valves 76a and 76b, liner 62, and bottle 52.

Pistons 78a and 78b move within their respective cylinders 72a and 72b compressing the formulation components within the cylinders as well as springs 84a and 84b. Balls 74a and 74b are firmly seated in valve seats 68a and 68b, operating as check valves to prevent the formulation components in the cylinders from being forced into respective reservoirs 54a and 54b. The formulation components are forced upwardly through the piston bores 80a and 80b in each piston respectively, the hydraulic pressure within each cylinder unseating balls 90a and 90b from valve seats 88a and 88b at the tops of the piston bores 80a and 80b. The formulation components are forced from the cylinders through the piston bores into ducts 86a

- 19 -

and 86b in the spout, the components being dispensed from dispensing nozzles 94a and 94b in the ducts and into the hand of the user who mixes the components to form the complete formulation, ready for use.

5        Upon release of spout 82, springs 84a and 84b push the pistons 78a, 78b to their biased position spaced away from check valves 76a and 76b. Upward motion of the pistons creates a suction within the cylinders 72a and 72b and the piston bores 80a and 80b. The suction seats balls  
10 90a and 90b on valve seats 88a and 88b to prevent formulation components in ducts 86a and 86b from being drawn back into the piston bores 80a and 80b. Simultaneously, the suction causes balls 74a and 74b to lift off their respective seats 68a and 68b as the  
15 formulation components are drawn up through dip tubes 66a and 66b from reservoirs 54a and 54b and into cylinders 72a and 72b. The pump is then ready to dispense the formulation components on the next actuation.

As explained above, each component of the formulation  
20 is pumped from its reservoir by its own piston-cylinder combination. The amount of a particular component dispensed from the pump is proportional to the volume of the cylinder swept by its piston, which pumps that particular component. Since the strokes of both pistons  
25 are equal, the relative amount of component dispensed from one reservoir in relation to the other reservoir is determined by the ratio of the cylinder diameters. Cylinders having equal diameters will have equal swept

- 20 -

volumes and, thus, dispense equal amounts of formulation component from each reservoir. If the diameter of one of the cylinders is made smaller relative to the other cylinder, then the piston associated with the smaller cylinder will sweep a smaller volume, and a smaller amount of component will be dispensed from the respective reservoir. Thus, the pump can be designed to automatically dispense formulation components in the proper relative proportions for maximum effectiveness simply by having cylinders with different diameters, the ratio of the diameters being proportional to the desired ratio of formulation components to be dispensed.

The package embodiments of the present invention are designed to be used by the lay person who has an acne condition or a lay person who applies the composition to the individual with the condition. Accordingly, the packages are designed so that the user is able to dispense on a consistent basis effective amounts of the active ingredients comprising the pharmaceutical composition.

Thus, the need for a pharmacist to prepare the composition is eliminated. The specific amount of each of the active ingredients present in the package will depend on the identity of the active ingredient, taking into account its known effect on treatment of the acne condition and the proportion of the ingredients that are desired in the composition that is formed by mixing the active ingredients. Also, the package may contain an optimal dosage of the active ingredients of the composition by



- 21 -

including in each packet the appropriate volume and concentration of the active ingredient. For example, in the use of a peroxide/antibiotic composition, it may be desired that the peroxide comprises a greater proportion  
5 of the composition than the antibiotic.

It will be appreciated that the packets of the single-dose package, as described herein, contain relatively small volumes of the components from which the composition is formulated. For example, an embodiment of  
10 such a packet contains about 0.425 g of one component and about 0.425 g of a second component. The dispensed components are within plus or minus 0.025 g of the 0.425 g in the packet, allowing for about a 6% tolerance. The filling of the packets with effective dosage amounts of  
15 the components is essential to providing an effective acne-treating composition. This is difficult to accomplish with components that are in a form of the type described herein, for example, in gel form. In order to fill the packets with the accurate dosages of components,  
20 it has been found that it is important to rid the components of air trapped within the components. To avoid trapped air, the components are formulated preferably using a process which avoids the introduction of air into the components and removes trapped air from the  
25 components. This is accomplished by preparing the components under a vacuum, for example, about -15 to about -30 psi. By substantially eliminating trapped air from the components, it is possible to fill the packets or

- 22 -

other appropriate containers with a high degree of accuracy. This, in turn, allows the lay person to prepare the acne-treating composition within the desired concentration parameters.

5       The form of the composition that is applied to the skin should be such that it adheres to the skin for a sufficient period of time to allow the active ingredients to act on the affected area. As an example, for  
embodiments which include a peroxide and an antibiotic,  
10 the composition will be in a form which adheres to the skin for a sufficient period of time for the peroxide to inactivate extracellular lipases and inhibit the formation of free fatty acids in the skin and for the antibiotic to reduce the concentration of acne-associated bacteria in  
15 the affected area. For this purpose, the acne-treating composition should have a viscosity such that the composition adheres to the skin. This requires that, when the components that make up the composition are combined together, they form a composition which has a viscosity  
20 such that the composition adheres to the skin. The forms of the components and the viscosities thereof can be different and vary substantially, provided that they can be combined to form a composition having the desired viscosity characteristics. For example, the components  
25 may be in forms ranging from liquids to solids. In preferred embodiments, as described in detail below, the component is in a form which has properties intermediate the liquid and solid states. Examples of such forms

- 23 -

include gels, lotions, creams, salves, and ointments. In a particularly preferred embodiment, the components are in gel form.

The package of the present invention, as exemplified in Figures 1 and 2, includes the separated components in dosages which are effective in forming, upon mixing, an effective acne-treating composition. In preferred form, the package is designed and the components are formulated to enable the user to dispense on a consistent basis substantially all of the contents of the package. Accordingly, the contents are made available for combining into an acne-treating composition in which the components are present in the desired dosage. The preferred means of accomplishing this is to formulate components which have viscosities that enable the user to dispense the components from the package without undue effort. It is preferred also that the components be in the same form. For example, if one component is in gel form, it is preferred that the other component be in gel form. In addition, it is preferred that the components have similar viscosities. The term "similar viscosity" means that when a uniform "dispensing" force is applied to the separated components, substantially equal amounts of the components are dispensed simultaneously from their respective chambers. In a particularly preferred form, the components have viscosities, as measured by 1V-101 Brookfield Helipath T-F @ 1.5 rpm that are within about 50%, preferably within about 25% of each other.

- 24 -

Standard procedures known in the art may be used to determine viscosity. To perform the viscosity test, a test sample was transferred to a 1lb jar (89 mm in diameter and 92 mm in height) while tapping on the jar to eliminate air. After the transfer, the jar was left to sit for one hour at 25.0°C. A Brookfield Model LV series rotational viscometer equipped with a helipath and a T-F spindle was used to measure the viscosity of the sample. The spindle of the viscometer was placed in the sample and the height of the spindle was adjusted so that the crosspiece of the spindle was covered by about 1/4" of the sample. The rotational speed of the spindle was set to 1.5 rpm. The helipath and the viscometer were simultaneously started. After one minute, the needle lock was engaged. When the needle was displayed in the window, the viscometer was shut off. A reading was taken and recorded on the 0-100 scale. Additional tests may be performed to obtain additional readings. If additional tests are performed, the viscometer rotation should be resumed and at least one rotation allowed prior to the release of the needle lock. Timing of the additional sample readings is to be commenced from the time of the release of the needle lock.

In particularly preferred embodiments, each of the components is a gel which has a viscosity of about 200,000 cps to about 500,000 cps.

There follows a description of exemplary ingredients comprising the acne-treating composition of the present

- 25 -

invention, including a description of exemplary pharmaceutically active ingredients and exemplary ingredients that are combined therewith to place the composition in a suitable form for application to the face  
5 and for dispensing the components of the composition from the package of the present invention.

In the discussion which follows, the amount of an ingredient in the acne-treating composition is expressed in weight percent based on the total weight of the  
10 composition (% w/w - composition). The amount of an ingredient in one of the components is expressed in weight percent based on the total weight of the component (% w/w - component).

The acne-treating composition includes an antibiotic  
15 which is effective against acne-associated bacterial species. An antibiotic is effective against acne-associated bacterial species if it kills or inhibits the growth of a bacterial population. The term "acne-associated bacterial species" refers to bacterial species  
20 found at the site of an acne lesion, in particular *Propionibacterium acnes* (*P. acnes*).

Exemplary antibiotics include lincomycin, clindamycin (a lincomycin derivative), and erythromycin and their pharmaceutically acceptable salts and esters such as their  
25 hydrochlorides and phosphates.

Lincomycin is a derivative of the amino acid trans-L-4- $\alpha$ -propyl-hydrinic acid coupled to a derivative of an octose substituted by a methylmercaptyl group. Lincomycin

- 26 -

antibiotics are described in U.S. Patent Nos. 3,475,407; 3,509,127; 3,544,551 and 3,513,155.

Clindamycin is the 7-deoxy,7-chloro derivative of lincomycin, and is otherwise known as methyl 7-chloro-  
5 6,7,8, trideoxy-6-[[ (1-methyl-4-propyl-2-pyrrolidinyl) carbonyl] amino]-1-thio-L-threo- $\alpha$ -D-galacto-octopyranoside. In the use of clindamycin as the antibiotic, it is recommended that there be used clindamycin phosphate which is converted *in vivo* to active clindamycin.

10 A preferred antibiotic for use in the present invention is erythromycin, which is a macrolide antibiotic produced from a strain of *Saccharopolyspora erythraea* (formerly *Streptomyces erythreus*). It is a base and readily forms salts with acids. Erythromycin is otherwise  
15 known as [(3R\*, 4S\*, 5S\*, 6R\*, 7R\*, 9R\*, 11R\*, 12R\*, 13S\*, 14R\*)-4-[(2,6-dideoxy-3-C-methyl-3-O-methyl- $\alpha$ -L-ribo-hexopyranosyl)-oxy]-14-ethyl-7,12,13-trihydroxy-3,5,7,9,11,13-hexamethyl-6-[[3,4,6-trideoxy-3-(dimethylamino)- $\beta$ -D-xylo-hexopyrano-  
20 syl]oxy]oxacyclotetradecane-2,10-dione].

The antibiotic is present in the composition in a pharmaceutically effective concentration. Such concentration will vary depending on the antibiotic used and the nature of the individual being treated. For  
25 guideline purposes, it is recommended that the antibiotic be present at a concentration of about 0.5 to about 6% w/w - composition, preferably about 0.8 to about 5% w/w - composition. In particularly preferred embodiments of the

- 27 -

invention utilizing erythromycin, the erythromycin is at a concentration of about 1.8 to about 4.5% w/w - composition. In preferred embodiments utilizing clindamycin, the clindamycin is at a concentration of  
5 about 0.7 to about 2% w/w - composition.

In addition to the antibiotic, the package of the present invention can include any other active ingredient that is effective in treating acne and that may be combined with an antibiotic to form a composition which  
10 tends to degrade prematurely. Examples of such active ingredients include corticosteroids, tretinoin, antifungal compounds, alpha-hydroxy acids and beta-hydroxy acids.

In preferred embodiments, the acne-treating composition preferably includes an antibiotic as described  
15 above and an oxidizing agent which inhibits the formation of free fatty acids in the skin primarily through inactivation of extracellular lipase enzymes. Preferably, the composition includes a "peroxide" oxidizing agent. The term "peroxide" means an organic compound containing  
20 an oxygen-oxygen bond capable of cleaving and forming oxygen free-radicals. A variety of peroxides may be used in the practice of the present invention. The peroxides include peroxyacids of carboxylic acids, peroxyesters of carboxylic acids and the dimeric product of carboxylic  
25 peroxyacids. Exemplary peroxides include t-butyl peroxyesters of straight and branched chain aliphatic carboxylic acids, and dimeric peroxides such as lauroyl peroxide and benzoyl peroxide.

- 28 -

In preferred embodiments utilizing a peroxide as the oxidizing agent, the peroxide should be of high purity and in the form of finely divided crystalline particles, preferably micronized particles having a mean average  
5 particle size of less than 35 microns. A preferred peroxide for use in the present invention is benzoyl peroxide, and the most preferred is micronized benzoyl peroxide 70% hydrous.

The oxidizing agent should be present in the  
10 composition in a pharmaceutically effective concentration. Such concentration will vary depending on the type of oxidizing agent used and the nature of the individual being treated. For guideline purposes, it is recommended that the composition contain about 1% to about 30% w/w -  
15 composition of oxidizing agent and preferably about 2.5% to about 15% w/w - composition oxidizing agent. In especially preferred embodiments, the oxidizing agent is at a concentration of about 3 to about 6% w/w - composition.

20 Active ingredients which are used in formulating the composition of the present invention and which are normally in solid form are advantageously converted into a liquid form in the preparation of the components from which the composition is made. For example, to prepare a  
25 liquid form of an active ingredient, it is combined with an appropriate solvent, the identity of which will depend on the particular active ingredient used. For example, if the active ingredient is highly soluble in water, then



- 29 -

water may be used as the solvent. Water may also be added to the composition as a diluent. If the active ingredient has low solubility in water, another appropriate solvent such as an alcohol may be used as the solvent.

5 Any pharmaceutically acceptable and suitable material can be used to impart to the component the desired viscosity. Viscosity-enhancing agents are well known. Preferably, a gelling agent is used. The gelling agent may be selected both as to type and quantity to give  
10 products of desired viscosities.

A variety of gelling agents may be used in the practice of the invention. Preferred gelling agents are pure micro-crystalline cellulose, colloidal magnesium-aluminum silicate, hydroxypropyl methyl cellulose and the  
15 so-called hydroxylated vinylic polymers, particularly, those disclosed in U.S. Patent No. 2,798,053. Preferred hydroxylated vinylic polymers are those described generally as interpolymers of a monomeric monolefinic acrylic acid and from about 0.1% to about 10% by weight  
20 based on the total monomer of a monomeric polyether of an oligosaccharide in which the hydroxyl groups that are modified are esterified with allyl groups, with said polyether containing at least two allyl ether groups per oligosaccharide molecule. Commercially available  
25 interpolymers of this type are known as "carbomers" marketed under the trademark Carbopol®. Such interpolymers are described as being polymers of acrylic acid cross-linked with about 1% of a polyallyl ether of

- 30 -

sucrose having an average of about 5.8 allyl groups for each sucrose molecule. These polymers have molecular weight in the order of magnitude of about 1,000,000. Such polymers are available from the B.F. Goodrich Chemical  
5 Company and are sold under such trademarks as Carbopol® 934, Carbopol® 940, and Carbopol® 941. A particularly preferred Carbopol is Carbopol® 934 NF.

The gelling agent should be present in the composition at a concentration that provides sufficient  
10 viscosity to the composition to allow the composition to adhere to the skin for a sufficient period of time to allow the active ingredients to act on the affected areas. The effective concentration will depend on the particular gelling agent used and the nature of the component to be  
15 gelled. In some embodiments, two or more gelling agents will be present. For guideline purposes, it is recommended that the gelling agent(s) comprise about 1 to about 4% w/w - composition, preferably about 1.5 to about 3% w/w - composition.

20 In a highly preferred composition of the present invention, hydroxypropylcellulose is used as a gelling agent.

In a preferred embodiment, a carbomer such as the Carbopol® interpolymers described hereinabove are used as a  
25 gelling agent in the oxidizing agent component and hydroxypropylcellulose is used as a gelling agent in the antibiotic component. Hydroxypropylcellulose is a propylene glycol ether of cellulose.

- 31 -

The preferred hydroxypropylcellulose used in the practice of the invention is available from Aqualon as Klucel® GF having a weight average molecular weight of 370,000.

5 A surface active agent may be included in the composition. The surface active agent functions as a processing aid which facilitates even distribution of the oxidizing agent (such as benzoyl peroxide) in the oxidizing agent component gel. The surface active agent may also  
10 function to retard the degradative interaction between an oxidizing agent and an antibiotic. Any pharmaceutically acceptable surface active agent may be used. An example of such a surface active agent is dioctyl sodium sulfosuccinate. The surface active agent is present at a concentration  
15 sufficient to facilitate even distribution of the oxidizing agent and optionally to retard the aforementioned degradative interaction. For guideline purposes, it is recommended that the surface active agent comprise about 0.05% to about 1% w/w - composition. In preferred embodiments, dioctyl sodium  
20 sulfosuccinate 75% is present at a concentration of about 0.07 to about 0.3% w/w - composition.

The preferred composition includes also a base. The base is added to neutralize the gelling agent (carbomer). The preferred pH range for the carbomer containing gels is about  
25 3 to about 7. For guideline purposes, the concentration is about 0.1 to about 1% w/w - composition, preferably about 0.15 to about 0.4% w/w - composition. Suitable bases include, but are not limited to, sodium hydroxide. Other

- 32 -

bases, including amines and hydroxides, may be used, for example, triethanolamine, PEG-15 cocamine and potassium hydroxide. In especially preferred embodiments, sodium hydroxide is used at a concentration of about 0.2 to about 5 0.35% w/w - composition.

Examples of other optional ingredients that may be present in an acne-treating composition are as follows. The composition can include a moistening agent (for example, propylene glycol) at a concentration that functions as a 10 solvent for methyl- and propylparabens. Preferably the moistening agent comprises about 0.5 to about 4% w/w - composition, most preferably about 1.5 to about 2.5% w/w - composition. The composition may include also an anti-microbial preservative (for example, methylparaben and 15 propylparaben) at a concentration sufficient to inhibit microbial growth in the composition. Preferably, the composition comprises about 0.01 to about 1% w/w - composition, most preferably about 0.03 to about 0.07% w/w - composition.

20 As mentioned above, a preferred composition for use in the practice of the present invention includes pharmaceutically effective amounts of benzoyl peroxide and erythromycin and hydroxypropylcellulose. A particularly preferred form of this composition comprises: (A) benzoyl 25 peroxide at a concentration of about 2.5 to about 15% w/w - composition, most preferably about 3 to about 6% w/w - composition; (B) erythromycin at a concentration of about 0.8 to about 5% w/w - composition, most preferably about 1.8 to

- 33 -

about 4.5% w/w - composition; and (C) hydroxypropylcellulose at a concentration of about 0.5 to about 4% w/w - composition, most preferably about 1 to about 3% w/w - composition.

5        Preferably the composition includes an additional gelling agent, preferably Carbopol® 934 at a concentration of about 0.5 to about 2% w/w - composition, most preferably about 0.7 to about 1.5% w/w - composition. In this particularly preferred form of the composition, water is used  
10 to dissolve the benzoyl peroxide at a concentration of about 35 to about 45% w/w - composition, most preferably about 42 to about 44% w/w - composition and an alcohol, preferably ethanol (SD #40B Alcohol as described in 27 CFR §21.76) is used to dissolve the erythromycin at a concentration of about  
15 40 to about 48% w/w - composition, most preferably about 43 to about 45% w/w - composition. Optional, but preferred ingredients, of this particularly preferred form of the composition comprise sodium dioctyl sulfosuccinate 75% at a concentration of about 0.05 to about 0.5% w/w - composition,  
20 most preferably about 0.08 to about 0.12% w/w - composition and sodium hydroxide at a concentration of about 0.05 to about 0.4% w/w - composition, most preferably about 0.1 to about 0.3% w/w - composition.

There follows a description of exemplary components  
25 which can be combined to form the acne-treating composition of the present invention.

Compositions for use in the practice of the present invention are prepared by combining at least two separate

- 34 -

active ingredient-containing components. Preferably, a first component used to prepare the composition contains an oxidizing agent and the second component contains an antibiotic. In preferred embodiments of the acne-treating composition, the oxidizing agent is present in equal or greater amounts than the antibiotic, preferably the ratio of oxidizing agent to antibiotic being about 1:1 to about 30:1. In highly preferred embodiments, the ratio of oxidizing agent to antibiotic is about 1:1 to about 5:1.

10 In preferred embodiments of the present invention, the antibiotic component comprises clindamycin or erythromycin, a solvent for the antibiotic, and a gelling agent. Additional ingredients which may be present include moistening agents, anti-microbial agents and a base.

15 In a preferred antibiotic component containing clindamycin ("clindamycin component"), the clindamycin is in the form of clindamycin phosphate and is present in a concentration such that the acne-treating composition includes a pharmaceutically effective concentration of the clindamycin phosphate. A milligram of clindamycin phosphate contains approximately 841µg of clindamycin. Accordingly, a 0.7-2% w/w-composition clindamycin will require approximately 0.8-2.3% w/w-composition clindamycin phosphate. Preferably the clindamycin phosphate concentration comprises about 1.5 20 to about 3.5% w/w - component. The clindamycin component includes additionally a solvent for the clindamycin. In preferred embodiments, the solvent is purified water which comprises about 88 to about 93% w/w - component. The

- 35 -

clindamycin component also includes a gelling agent which imparts thereto a viscosity (about 300,000 to about 500,000 cp) which is similar to the viscosity of the other component used in the composition (preferably a benzoyl peroxide-containing component). The preferred gelling agent is Carbopol® 934 which comprises about 1 to about 2% w/w - component. The clindamycin component can also include a base for the purpose of neutralizing the carbomer (Carbopol® 934) gelling agent. The desired pH range for the carbomer gels is about 3 to about 7. Preferably the base is sodium hydroxide in a concentration of about 0.1 to about 0.3 w/w - component. Other preferred ingredients in the clindamycin component are: (A) the moistening agent propylene glycol at a concentration of about 2 to about 6% w/w - component; and (B) the antimicrobial agents methylparaben and propylparaben at concentrations of about 0.04 to about 0.2% w/w - component.

In a preferred antibiotic component containing erythromycin, the erythromycin is present in a concentration such that the acne-treating composition includes a pharmaceutically effective concentration of the erythromycin. Preferably the erythromycin concentration comprises about 5.5 to about 8.5% w/w - component. As erythromycin has limited solubility in water, the erythromycin is dissolved in alcohol, preferably SD alcohol #40-B 190 proof, in a concentration of about 87 to about 92% w/w - component. The erythromycin component also includes a gelling agent which imparts to the component a viscosity (about 200,000 cps to about 400,000 cps) that is similar to the viscosity of the

- 36 -

other component used in the composition (preferably a benzoyl peroxide-containing component). Hydroxypropylcellulose is the preferred gelling agent and is present preferably in a concentration of about 3 to about 5% w/w - component.

5 A preferred oxidizing agent component comprises benzoyl peroxide (for example, 70% hydrous) in a concentration such that the acne-treating composition contains a pharmaceutically effective concentration of the benzoyl peroxide. Preferably the benzoyl peroxide comprises about 8  
10 to about 12% w/w - component. The benzoyl peroxide is preferably dissolved in water at a concentration of about 80 to about 85% w/w - component. The oxidizing agent component also includes a gelling agent which imparts to the component a viscosity (about 200,000 to about 500,000 cp) which is  
15 similar to the viscosity of the other component used to prepare the acne-treating composition (preferably an antibiotic component). A preferred gelling agent is Carbopol® 934 at a concentration of about 1 to about 3% w/w - component. The preferred oxidizing agent component comprises  
20 also the surface active agent dioctyl sodium sulfosuccinate at a concentration of about 0.1 to 0.3% w/w - component. Another preferred ingredient of the oxidizing agent component is sodium hydroxide which is present at a concentration of about 0.3 to 0.5% w/w - component.

25

### Examples

The following examples are illustrative of the present invention. The term "USP" as used herein refers to chemicals



- 37 -

that conform to the tolerances set forth in the United States Pharmacopeia/National Formulary. (The term "NF" refers to National Formulary standards.) The United States Pharmacopeia/National Formulary is a compendium of pharmaceutical formulations widely used as a standard reference. The term "USP" refers to United States Pharmacopeia standards. The USP/NF is maintained by the United States Pharmacopeial Convention, Inc. The USP24/NF19 was issued January, 2000.

10 Example 1 - Preparation of Benzoyl Peroxide Gel/Erythromycin Gel Package

(A) Benzoyl Peroxide Gel

A benzoyl peroxide gel containing the following ingredients was prepared as follows.

15	<u>Ingredients</u>	<u>Weight Percent</u>
	Purified Water, USP	82.495
	Carbopol® 934, NF	1.900
	Sodium Hydroxide, NF	0.405
	Benzoyl Peroxide, USP 70%	15.000
20	Diocetyl Sodium Sulfosuccinate 75%	0.200

Purified water was added to a tank (hereinafter, the "Phase A tank") and sodium hydroxide pellets were added to the water. The contents of the tank were mixed until the pellets fully dissolved in the water and the solution cooled to 22-30 °C.

25. Purified water was introduced into a second tank (hereinafter, the "main tank") and the water was cooled to 22-25°C. Carbopol® 934 was dispensed into the main tank and

- 38 -

the tank mixed. The contents of the Phase A tank were slowly added to the contents of the main tank while continuously mixing the contents of the main tank at a temperature of 22-30°C.

5        Purified water was added to a third tank (hereinafter, the "phase B tank"). The dioctyl sodium sulfosuccinate was added to the tank, followed by addition of benzoyl peroxide. The contents of the tank were mixed producing a slurry. The Phase B tank was connected to the main tank and the benzoyl  
10 peroxide slurry was pumped into the main tank.

The contents of the main tank were mixed under a 10-15" psi vacuum and transferred to clean and sanitized bulk storage containers. The resultant gel had a viscosity of 360,000 cps. The viscosity was tested using method 1V-101  
15 Brookfield helipath T-F @ 1.5 rpm.

- 39 -

(B) Erythromycin Gel

An erythromycin gel containing the following ingredients was prepared as follows.

	<u>Ingredients</u>	<u>Weight Percent</u>
5	S.D. Alcohol #40-B 190 Proof	89.10
	Erythromycin, USP	6.90
	Hydroxypropylcellulose, NF	4.00

Specially denatured (SD) Alcohol #40-B, 190 proof was added to a vessel. Erythromycin was added while mixing.

10 Mixing was continued until the erythromycin completely dissolved. Hydroxypropylcellulose (Klucel® GF manufactured by Aqualon) was then added while mixing. Mixing was performed at -15 to -30 psi to facilitate deaeration. Mixing may be performed in an open or closed vessel. For commercial

15 preparations, mixing will be performed in a closed vessel. After mixing, the tank was sealed and held for a minimum of 12 hours at room temperature in order to allow any entrapped air to dissipate and to assure that all of the hydroxypropylcellulose had gelled. Any evaporated alcohol

20 was replaced with SD Alcohol #40-B to the target weight and the mixture was mixed at -15 to -30 psi. Mixing may be performed in an open or closed vessel. For commercial preparations, mixing will be performed in a closed vessel. The mixture was then transferred into clean and sanitized

25 steel drums. The resultant gel had a viscosity of 360,000 cps. The viscosity was tested using method 1V-101 Brookfield helipath T-F @ 1.5 rpm.

(C) Packaging the Gels

The packaging is performed using a Bartelt horizontal, form-fill-and-seal machine.

The benzoyl peroxide and the erythromycin gels were transferred to pressure vessels charged by inert gas. A pressure regulator was used to keep the pressure level constant. The pressure level was set up to 80 psi for benzoyl peroxide and up to 40 psi for erythromycin gel. A vertical linear piston-metering chamber was installed in each of the gel filling lines. The volume of gel in the chamber was kept constant by the pressurized vessels continuously forcing gel into the metering chamber which assures filling precision and accuracy of viscous gels at high speeds and low volumes. The stroke of the piston in the metering chamber is initiated by a programmable logic controller (PLC) signal synchronized with the movement of the rest of the filling line. The piston drives the gels to a filling pump. A horizontal linear positive displacement micro-pump (*Hi-Bar pump*), with a packet-dosing chamber, was installed at the terminal end of the filling line. The dosing chamber was filled by the action of the metering chamber piston. The dosing chamber feeds a loop of clear tubing terminating in a vertical piston, at the filling station.

The gels are packaged in a two-packet laminated foil package containing the benzoyl peroxide gel in one packet and the erythromycin gel in the other packet. The benzoyl peroxide gel (0.425g) was filled into one packet while the erythromycin gel (0.425g) was simultaneously filled into the

second packet. The dispensed gels are within plus or minus 0.025g of the target weight of 0.425g, allowing for a 6% tolerance. After each packet was filled and sealed, the packets were folded and cold-sealed together using a glue  
5 adhesive. The packets are constructed of a laminated foil consisting of an inner barrier of low-density polyethylene; a tie layer of ethylene/acrylic acid copolymer; a main barrier of foil; a tie layer of ethylene/acrylic acid copolymer; and an outer barrier of polyester.

10 Combining the benzoyl peroxide gel with the erythromycin gel provides an acne-treating composition having the following ingredients by weight%.

INGREDIENT	% W/W COMPOSITION
Purified Water, USP	43.50
15 Carbopol® 934, NF	0.95
Sodium Hydroxide, NF	0.20
Benzoyl Peroxide, 70% Hydrous, USP	5.25^
Diocetyl Sodium Sulfosuccinate, DF	0.10
SD Alcohol #40-B/190°, DF	44.55
20 Erythromycin, USP	3.45
Hydroxypropylcellulose, NF	2.00

Example 2 - Preparation of Benzoyl Peroxide/Clindamycin Gel Package

The benzoyl peroxide gel is prepared as described in

25 Example 1.

A clindamycin gel containing the following ingredients was prepared as follows.

	<u>Ingredients</u>	<u>Weight Percent</u>
	Purified Water, USP	91.28
5	Carbomer 934	1.50
	Sodium Hydroxide, NF	0.27
	Propylene Glycol, USP	4.00
	Methylparaben, NF	0.14
	Propylparaben, NF	0.06
10	Clindamycin Phosphate, USP	2.75

Purified water was added to a tank (hereinafter, the "Phase A tank") and sodium hydroxide pellets were added to the purified water and the contents of the tank were mixed until the pellets fully dissolved in the purified water and  
15 the solution cooled to 22-30°C.

Purified water was added into a second tank (hereinafter, the "main tank") and Carbomer 934 was dispensed into the main tank. The contents of the phase A tank were slowly added to the contents of the main tank while  
20 continuously mixing the contents of the main tank and deaerating under a vacuum. The temperature was maintained at 22-30°C.

Purified water was added to a third tank (hereinafter, the "phase B tank"). Clindamycin phosphate was added and  
25 mixed until dissolved. Once the clindamycin phosphate was completely dissolved, the contents of the tank were pumped into the main tank and mixed under vacuum.

Purified water and propylene glycol were added to a fourth tank. The tank was heated to 35-40°C. Methylparaben

was then added and the contents of the tank were mixed until all the methylparaben was dissolved. Propylparaben was then added and the contents of the tank mixed until all the propylparaben was dissolved. The contents were transferred  
5 into the main tank.

While continuously mixing, a vacuum was turned on and the mixing continued until the material was homogenous. The contents of the tank were then transferred to stainless steel drums. The resultant gel had a viscosity of 370,000 cps.  
10 The viscosity was tested using method IV-101 Brookfield helipath T-P @ 1.5 rpm.

The benzoyl peroxide and the erythromycin gels were transferred to pressure vessels charged by inert gas. A pressure regulator was used to keep the pressure level  
15 constant. The pressure level was set up to 80 psi for benzoyl peroxide and up to 40 psi for erythromycin gel. A vertical linear piston-metering chamber was installed in each of the gel filling lines. The volume of gel in the chamber was kept constant by the pressurized vessels continuously  
20 forcing gel into the metering chamber which assures filling precision and accuracy of viscous gels at high speeds and low volumes. The stroke of the piston in the metering chamber is initiated by a PLC signal synchronized with the movement of the rest of the filling line. The piston drives the gels to  
25 a filling pump. A horizontal linear positive-displacement micro-pump, with a packet dosing chamber, was installed at the terminal end of the filling line. The dosing chamber was filled by the action of the metering chamber piston. The

dosing chamber feed a loop of clear tubing terminating in a vertical piston, at the filling station.

For the final packaging step, 0.425g of a 10% benzoyl peroxide gel was filled into one packet while 0.425g of a 2% clindamycin gel was simultaneously filled into the second packet. The dispensed gels are within plus or minus 0.025 g of the target weight of 0.425g, allowing for a 6% tolerance. After each pouch was filled and sealed, the pouches were folded and cold-sealed together.

Combining the benzoyl peroxide gel with the clindamycin gel provides an acne-treating composition having the following ingredients by weight%.

	INGREDIENT	% W/W COMPOSITION
	Purified Water, USP	89.14
15	Carbomer 934, NF	1.70
	Sodium Hydroxide, NF	0.34
	Propylene Glycol, USP	2.00
	Diethyl Sodium Sulfosuccinate 75%, DF	0.10
	Methylparaben, NF	0.07
20	Propylparaben, NF	0.03
	*Clindamycin Phosphate, USP	1.37
	Hydrous Benzoyl Peroxide, USP	5.25

\*The % w/w for the clindamycin phosphate is based solely on % w/w. The active substance in the product is clindamycin. Clindamycin phosphate is converted *in vitro* to active clindamycin. The actual amount of clindamycin as clindamycin phosphate is based on several factors including activity. The actual amount of material added is based on 6.875 kg of clindamycin phosphate.



Claims

1. A package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, said composition tending to degrade prematurely, one of said components comprising an oxidizing agent and another of said components comprising an antibiotic which is effective against acne-associated bacterial species, said components separated one from the other in said package, said one component having a viscosity within about 50% of the viscosity of said other component.

2. The package of Claim 1 wherein said component comprising an oxidizing agent comprises about 8 to about 12% w/w - component oxidizing agent.

3. The package of Claim 1 wherein said component comprising an antibiotic comprises about 1.5 to about 8.5% w/w - component antibiotic.

4. The package of Claim 1 wherein the ratio of said oxidizing agent to said antibiotic is about 1:1 to about 30:1.

5. The package of Claim 4 wherein the ratio of said oxidizing agent to said antibiotic is about 1:1 to about 5:1.

6. The package of Claim 1 wherein said oxidizing agent is benzoyl peroxide.

7. The package of Claim 1 wherein said antibiotic is selected from the group consisting of erythromycin and clindamycin.

8. The package of Claim 7 wherein said antibiotic is erythromycin.

9. The package of Claim 1 wherein said package comprises a dual packet laminated foil package.

10. A package comprising components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of said components comprising a benzoyl peroxide gel and another of said components comprising a gel of erythromycin, said components separated one from the other, said benzoyl peroxide gel having a viscosity within about 50% of the viscosity of said erythromycin gel.

11. The package of Claim 10 wherein said benzoyl peroxide gel comprises about 8 to about 12% w/w - component benzoyl peroxide.

12. The package of Claim 11 wherein said benzoyl peroxide gel comprises about 10% w/w - component benzoyl peroxide.

13. The package of Claim 10 wherein said erythromycin gel comprises about 5.5 to about 8.5% w/w - component erythromycin.

14. The package of Claim 13 wherein said erythromycin gel comprises about 7% w/w - component erythromycin.

15. The package of Claim 14 wherein the ratio of said benzoyl peroxide to said erythromycin is about 1:1 to about 30:1.

16. The package of Claim 15 wherein the ratio of said benzoyl peroxide to said erythromycin is about 1:1 to about 5:1.

17. The package of Claim 10 wherein said package is a dual packet laminated foil package.

18. A package comprising: (A) components which, upon being mixed, are capable of forming a pharmaceutical composition that is effective in treating acne, one of said components comprising a benzoyl peroxide gel and another of said components comprising a gel of erythromycin and hydroxypropylcellulose; and (B) containers for holding said

components in the package separated one from the other; said components having viscosities such that, upon the application of a uniform force to said components, substantially equal volumes of said components are capable of being dispensed simultaneously from said containers.

19. The package of Claim 18 wherein said benzoyl peroxide gel comprises about 8 to about 12% w/w - component benzoyl peroxide.

20. The package of Claim 18 wherein said erythromycin and hydroxypropylcellulose gel comprises about 5.5 to about 8.5% w/w - component erythromycin and about 3 to about 5% w/w - component hydroxypropylcellulose.

21. The package of Claim 18 wherein said benzoyl peroxide gel comprises about 10.5% w/w - component benzoyl peroxide and said gel of erythromycin and hydroxypropylcellulose comprises about 7% w/w - component erythromycin and about 4% w/w - component hydroxypropylcellulose.

22. The package of Claim 21 wherein said benzoyl peroxide gel has a viscosity from about 200,000 cps to about 500,000 cps and said erythromycin and hydroxypropylcellulose gel has a viscosity of from about 200,000 cps to about 400,000 cps.

23. A composition for the treatment of acne comprising:

- (a) benzoyl peroxide;
- (b) erythromycin; and
- (c) hydroxypropylcellulose.

24. The composition of Claim 23 including an additional gelling agent.

25. The composition of Claim 23 including a surface active agent.

26. The composition of Claim 23 including a base which increases the pH of said composition to a value at which said composition has an acceptable viscosity.

27. The composition of Claim 23 including an alcohol solvent for said erythromycin.

28. The composition of Claim 23 including a diluent.

29. A composition for the treatment of acne comprising:  
(a) benzoyl peroxide; (b) erythromycin; (c) hydroxypropylcellulose; (d) a gelling agent; (e) a surface active agent; (f) a base; (g) an alcohol; and (h) water.

30. The composition of Claim 29 wherein said gelling agent is a hydroxylated vinylic polymer.

31. The composition of Claim 29 wherein said surface active agent is dioctyl sodium sulfosuccinate.

32. The composition of Claim 29 wherein said base is sodium hydroxide.

33. The composition of Claim 29 wherein said alcohol is specially denatured ethanol.

34. A composition for the treatment of acne comprising about 5.25% w/w - composition benzoyl peroxide, about 3.45% w/w - composition erythromycin, and about 2% w/w - composition hydroxypropylcellulose.

35. The composition of Claim 34 comprising about 0.95% w/w - composition of a hydroxylated vinylic polymer, about 0.10% w/w - composition of dioctyl sodium sulfosuccinate, about 0.2% w/w - composition of sodium hydroxide, about 44.55% w/w - composition specially denatured ethanol and about 43.5% w/w - composition of water.

36. A package containing a first acne-treating composition and a second acne-treating composition, said compositions tending to degrade chemically upon being mixed with one another, said package comprising a first chamber containing said first composition and a second chamber containing said second composition, each of said chambers

having a sealed dispensing orifice, said orifices positioned adjacent to one another.

37. A package containing a first acne-treating composition and a second acne-treating composition, said compositions tending to degrade chemically upon being mixed with one another, said package comprising a first and second packet, said packets being attached to each other and each of said packets having a sealed dispensing orifice, said sealed dispensing orifices being positioned adjacent to each other so that, as said compositions are dispensed through the orifices in unsealed form, they are combined with each other.

38. A package containing a first acne-treating composition and a second acne-treating composition, said compositions tending to degrade chemically upon being mixed with one another, said package comprising a first and second packet, each of said packets having a top which includes a dispensing orifice that has a tear-off tab seal, a bottom, two sides in a parallel relationship to one another, and an exterior face and an interior face, said first packet containing said first composition and said second packet containing said second composition, said interior faces of said packets adjacent to and facing one another such that said tear-off tabs can be simultaneously torn off such that said compositions are dispensed and combined with each other as they exit said packets through said dispensing orifices.

1/4




FIG. 1




FIG. 2






FIG. 3




FIG. 4

4 / 4




FIG. 5

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC 7 A61K9/06 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, CHEM ABS Data

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 088 717 A (RORER INTERNATIONAL) 16 June 1982 (1982-06-16) claims 1-8,11,16-20	1-19, 36-38
Y	page 3, line 27 - line 47 examples 3-6	20-35
Y	US 4 954 487 A (EUGENE R. COOPER; ET AL.) 4 September 1990 (1990-09-04) examples 17,18	20-35
X	WO 99 02133 A (GIST BROCADES) 21 January 1999 (1999-01-21) claims 1-6 page 3, line 17 - line 24 page 4, line 17 - line 21 page 4, line 34 - line 37 page 6, line 9 - line 21 example 1	1-19, 36-38
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

13 November 2001

Date of mailing of the international search report

21/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

## INTERNATIONAL SEARCH REPORT

PCT/US 01/18033

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 93 15726 A (BAROODY, LLOYD; ET AL.) - 19 August 1993 (1993-08-19) claims 1,2 examples 1-13 ---	1-19, 36-38
X	CA 2 015 111 A (WOO, TOM; ET AL.) 23 October 1991 (1991-10-23) claims 1,4,5,11,13,72-79 -----	1-19, 36-38

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2088717	A	16-06-1982	AR 229405 A1	15-08-1983
			AT 381452 B	27-10-1986
			AT 285581 A	15-03-1986
			AU 557147 B2	11-12-1986
			AU 7202981 A	17-06-1982
			BE 889327 A1	16-10-1981
			CA 1179599 A1	18-12-1984
			CH 656307 A5	30-06-1986
			CH 647147 A5	15-01-1985
			DE 3124698 A1	08-07-1982
			DK 273981 A	09-09-1982
			ES 503797 D0	01-03-1983
			ES 8303087 A1	01-05-1983
			FR 2495471 A1	11-06-1982
			GB 2090135 A	07-07-1982
			IE 52227 B1	19-08-1987
			IT 1210608 B	14-09-1989
			JP 57099525 A	21-06-1982
			MX 9203750 A1	01-09-1992
			MX 6555 E	10-07-1985
			NL 8102997 A	01-07-1982
			NZ 197491 A	28-02-1985
			NZ 209389 A	28-02-1985
			PH 19245 A	14-02-1986
			SE 8103925 A	09-06-1982
			US 4497794 A	05-02-1985
			US 4692329 A	08-09-1987
			ZA 8104215 A	23-02-1983
US 4954487	A	04-09-1990	EP 0013459 A2	23-07-1980
			JP 55120514 A	17-09-1980
			AU 544969 B2	27-06-1985
			AU 7272081 A	14-01-1982
			CA 1165240 A1	10-04-1984
			DE 3172516 D1	07-11-1985
			EP 0043738 A2	13-01-1982
			IE 51377 B1	10-12-1986
			JP 1737953 C	26-02-1993
			JP 4020886 B	07-04-1992
			JP 57081408 A	21-05-1982
			NZ 197668 A	16-03-1984
			ZA 8104650 A	28-07-1982
WO 9902133	A	21-01-1999	AU 9259598 A	08-02-1999
			BR 9810985 A	15-08-2000
			CN 1262616 T	09-08-2000
			WO 9902133 A2	21-01-1999
			EP 0996428 A2	03-05-2000
			JP 2001509477 T	24-07-2001
WO 9315726	A	19-08-1993	AT 197248 T	15-11-2000
			AU 3478893 A	03-09-1993
			BR 9305884 A	19-08-1997
			CA 2130341 A1	19-08-1993
			CN 1077887 A ,B	03-11-1993
			DE 69329617 D1	07-12-2000
			DE 69329617 T2	23-05-2001
			DK 626844 T3	22-01-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9315726	A	EP 0626844 A1	07-12-1994
		EP 0960618 A2	01-12-1999
		ES 2153377 T3	01-03-2001
		FI 943783 A	17-08-1994
		JP 7504168 T	11-05-1995
		MX 9300848 A1	31-08-1994
		NO 943045 A	17-08-1994
		NZ 249021 A	27-08-1996
		PL 171127 B1	28-03-1997
		PT 626844 T	30-03-2001
		RU 2122412 C1	27-11-1998
		SG 48195 A1	17-04-1998
		WO 9315726 A1	19-08-1993
		US 5733886 A	31-03-1998
		US 6117843 A	12-09-2000
CA 2015111	A	23-10-1991	CA 2015111 A1
			23-10-1991

**THIS PAGE BLANK (USPTO)**